

Biological

Established 1832

Medical  
Serials

Fifth Series, No. 8

# THE IRISH JOURNAL OF MEDICAL SCIENCE

FORMERLY

THE DUBLIN JOURNAL OF MEDICAL SCIENCE

THE OFFICIAL ORGAN OF THE ROYAL  
ACADEMY OF MEDICINE IN IRELAND

EDITED BY

A. K. HENRY, M.B., F.R.C.S.I.



CAHILL & CO., LTD., 40 LOWER ORMOND QUAY, DUBLIN,  
AND 43 NEW OXFORD ST., W.C.2.

*To whom all Communications for the Editor and Books for Review  
are to be addressed.*

LONDON: SIMPKIN, MARSHALL & CO. EDINBURGH: JAMES THIN.

AGENT IN AMERICA:

G. E. STECHERT & CO., 151 to 155 WEST 25TH ST., NEW YORK.

Annual Subscription, £1 5s.

Single No. 2/6 net.

PUBLISHED MONTHLY.

OCTOBER, 1922.

THE FRENCH NATURAL MINERAL WATER

# VICHY-CÉLESTINS

(Property of the FRENCH STATE)



FOR DISORDERS OF  
THE LIVER,  
GOUT, RHEUMATISM,  
DIABETES, GRAVEL, Etc.

---

NATURAL VICHY SALTS  
For Drinking and Baths.

VICHY DIGESTIVE PASTILLES  
prepared with Natural Vichy Salts.

---

CAUTION.—Each bottle from the STATE  
SPRINGS bears a neck label with the word  
"VICHY-ETAT" and the name of the  
SOLE AGENTS:—

**INGRAM & ROYLE, LIMITED**

Bangor Wharf, 45 Belvedere Road, London, S.E.

And at LIVERPOOL and BRISTOL.

*Samples Free to Members of the Medical Profession.*

# J. & A. CHURCHILL

Now Ready. 80 Illustrations. 15s. net. Postage 8d.

## Medical Ophthalmology

By R. FOSTER MOORE, O.B.E., F.R.C.S., Asst. Ophthal. Surgeon,  
St. Bart's Hospital.

*"Most excellently has he succeeded in his aim. The book is packed with information. In our opinion this book is one of the most useful to the physician and to the ophthalmic surgeon alike that has appeared in this country in recent years. The illustrations admirably illustrate the text."*

—BRIT. MED. JOUR.

## Williams' Minor Surgery and Bandaging

17th Edition. 239 Illustrations. 10s. 6d. net. Postage, 6d.

## STUDENTS' SYNOPSIS SERIES

### Back and Edwards' Surgery

12s. 6d. net. Postage, 6d.

### Johnston's Regional Anatomy

11 Illustrations. 12s. 6d. net. Postage 6d.

### Burnet's Materia Medica

With Notes on Prescription Writing. 4s. 6d. net. Postage 3d.

### Jameson and Marchant's Hygiene

18 Illustrations. 18s. net. Postage 7d.

### Roberts' Physiology

73 Illustrations. 15s. net. Postage 6d.

### Cow's Pharmacology

16 Illustrations. 7s. 6d. net. Postage 3d.

### Underwood's Dentistry

10 Illustrations. 9s. 6d. net. Postage 3d.

### Pearce Gould's Surgical Pathology

6s. net. Postage 4d.

### Surgical Diagnosis

By W. H. C. ROMANIS, F.R.C.S., Lecturer on Practical  
Surgery, and Assistant Surgeon, St. Thomas's Hospital.  
Nearly Ready. 8s. 6d. net. Postage 6d.

7th Edition. 31 Plates and 68 Text-figures. 21s. net. Postage 1s.

## Hewlett's Bacteriology

*"The work of a teacher of great experience. . . . The subject matter is well set out, and the diction is extremely clear and comprehensive, so that the book is perhaps the most readable of all the bacteriological text-books in English."*

—BRIT. MED. JOUR.

## Hewlett's Pathology

Fifth Edition. 18s. net. Postage 1s. Now Ready.

London: J. & A. Churchill, 7 Great Marlborough Street, W.1

# J. & A. CHURCHILL

## Gask and Wilson's Surgery

A Text-Book by Various Authors.  
With 39 Plates, 20 in Colour, and 467 Text-figures. 30s. net. Postage 1s. 3d.  
*"This book is a notable addition to our surgical literature."*—*The British Journal of Surgery.*

*"We are impressed not only with the completeness of this volume, but with the excellent balance of the articles by different writers. Each section is handled with the strength and accuracy of the expert."*—*The Lancet.*

## Jex-Blake's Physical Signs in the Chest and Abdomen

27 Illustrations. 9s. 6d. net. Postage 4d.

*"A concise and lucid introduction to an essential branch of medical practice."*—*The Lancet.*

## Latham and Torrens' Medical Diagnosis

With 74 Illustrations, some in Colours. 15s. net. Postage 1s.

*"A valuable handbook conveniently arranged, efficiently illustrated."*—*Edinburgh Medical Journal.*

## Panton's Clinical Pathology

13 Plates (11 Coloured) and 45 Text-figures. 14s. net. Postage 1s.

## Sequeira's Diseases of the Skin

3rd Edition. 52 Coloured Plates, 257 Text-figures. 38s. net. Postage 1s. 3d.

## Small's Text-Book of Botany

For Medical and Pharmaceutical Students.

With 1,350 Illustrations. 21s. net. Postage 1s.

## Taylor's Principles and Practice of Medical Jurisprudence

7th Edition. With 20 Engravings. 2 Vols. £3 3s. net. Postage 1s. 3d.

*"For the lawyer as well as the medical practitioner this standard work is invaluable. In every branch of law into which medical considerations enter it may always be consulted with profit."*—*The Justice of the Peace.*

## Taylor's Operative Surgery

With 300 Original Illustrations, many in Colour. 32s. net. Postage 1s. 3d.

*"Eminently practical . . . confidently recommended . . . a credit to the surgery of the kingdom . . . printing excellent."*—*The Lancet.*

## Hale-White's Materia Medica

17th Edition. 10s. 6d. net. Postage 5d.

## Whittaker's Surface Anatomy

3rd Edition. 17 Plates. 7s. 6d. net. Postage 6d.

## Eden and Lockyer's Gynaecology

2nd Edition. 24 Coloured Plates and 513 Text-figures. 36s. net. Postage 1s. 3d.

*"In every way the new edition maintains the high standard of its predecessor, and it will continue to rank as probably the most satisfactory, all-round text-book of moderate size on the subject published in this country."*—*British Medical Journal.*



# J. & A. CHURCHILL

## Starling's Physiology

3rd Edition. With 579 Illustrations. 25s. net. Postage 1s.

*"A monument to the masterly presentment of the subject by one of the most eminent of British physiologists."*—PHYSIOLOGICAL ABSTRACTS.

## Fleming's Short Practice of Medicine

3rd Edition. With 64 Illustrations. 21s. net. Postage 1s.

## Goodhart and Still's Diseases of Children

11th Edition. With 60 Illustrations. 32s. net. Postage 1s.

## Hartridge's Refraction of the Eye

16th Edition. 110 Illustrations. 7s. 6d. net. Postage 6d.

## Hartridge on the Ophthalmoscope

6th Edition. 4 Plates, 65 Text-figures. 6s. 6d. net. Postage 6d.

## Hawk's Practical Physiological Chemistry

7th Edition. 6 Colour Plates, 192 Text-figures (12 in Colour). 24s. net ;  
Postage 1s.

## Hewlett and Nankivell's Preventive Medicine

17 Illustrations. 18s. net. Postage 1s.

## Rowlands and Turner's Operations of Surgery

797 Illustrations, 40 in Colour. 2 Vols. £2 10s. net. Postage 1s. 6d.

*"The very best book of its kind in the British language."*

THE PRACTITIONER.

## Jellett's Short Practice of Midwifery

8th Edition. 207 Text-figures. 4 Coloured Plates. 18s. net. Postage 8d.

## Jellett's Practice of Gynaecology

4th Edition. 374 Illustrations. 11 Coloured Plates. 21s. net. Postage 1s.

## Wood Jones' Principles of Anatomy as seen in the Hand

With 2 Plates and 123 Text-figures. 15s. net. Postage 7d.

*"It can be said at once, and heartily, that the book is one worth reading, and eminently readable, well got up, with clear and explanatory illustrations, and could with advantage be placed in the hands of every medical student who has attained to that investigation and contemplation of his intermediate subjects."*—THE LANCET.

## Frazer's Anatomy

2nd Edition. With 219 Illustrations, many in Colour. 28s. net.  
Postage 1s.

*"The underlying principle of the book is sound and commendable, and the author has succeeded admirably in his aim."*—BRITISH MEDICAL JOURNAL.

London : J. & A. Churchill, 7 Great Marlborough Street, W.1

# J. & A. CHURCHILL

## Taylor's Medicine

Edited by E. P. POULTON, M.D., F.R.C.P., Physician to Guy's Hospital  
12th Edition. 24 Plates. 84 Text-Figures. 30s. net. Postage 1s.

*"The revision has been remarkably successful and complete, and as evidence of its up-to-date character attention may be drawn to the inclusion in the appendix of van den Bergh's tests for bilirubin in the blood serum, with a reference to Dr. W. J. McNee's contributions to the subject in our columns so recently as 13th May last. The diseases of metabolism and those of endocrine glands are considered together in a section prefaced by a short summary, which the Editor is so specially fitted to contribute, of basal metabolism; the account of diabetes mellitus follows and is naturally both full and up-to-date."*—BRITISH MEDICAL JOURNAL, 21st July, 1922.

## Bennett's Medical and Pharmaceutical Latin

New (3rd) Edition. 10s. 6d. net. Postage 5d.

## Berkeley and Bonney's Difficulties and Emergencies of Obstetric Practice

3rd Edition. 309 Illustrations. 36s. net. Postage 1s.

## Bowlby and Andrewes' Surgical Pathology

7th Edition. With 200 Illustrations. 30s. net. Postage 1s.

## Box's Post-Mortem Manual

A Handbook of Morbid Anatomy and Post-mortem Technique.

By CHARLES R. BOX, M.D., B.S.Lond., F.R.C.P., F.R.C.S.

New (2nd) Edition. With 22 Illustrations. 10s. 6d. net. Postage 5d.

## Burns' Introduction to Biophysics

With Foreword by D. NOEL PATON, M.D., LL.D., F.R.S.

With 85 Illustrations. 21s. net. Postage 9d.

## Corbin and Stewart's Physics and Chemistry

5th Edition. 200 Illustrations. 12s. 6d. net. Postage 6d.

## Craig's Psychological Medicine

3rd Edition. 27 Plates. 18s. net. Postage 1s.

## Craig's Nerve Exhaustion

6s. net. Postage 4d.

## Crowther's Radiography

By J. A. CROWTHER, M.A., SC.D., F.Inst.P., University Lecturer in Physics applied to Medical Radiology, Cambridge.

Now Ready. With 55 Illustrations. 7s. 6d. net. Postage 5d.

## Cushny's Pharmacology and Therapeutics

7th Edition. With 71 Illustrations. 24s. net. Postage 1s.

## Eden's Midwifery

5th Edition. 5 Plates (4 in Colour), 269 Text-figures. 24s. net. Postage 1s.

*"We have nothing but praise for this book."*—THE LANCET.

London: J. & A. Churchill, 7 Great Marlborough Street, W.I.

# LEWIS'S PUBLICATIONS.

TWELFTH Edition. Thoroughly Revised. With 9 Coloured Plates and 263 other illustrations. Demy 8vo. 24s. net; Postage 1s.

## SWANZY'S HANDBOOK OF DISEASES OF THE EYE AND THEIR TREATMENT

Revised and Edited by LOUIS WERNER, M.B. F.R.C.S.I., Sen. Mod., Univ. Dub.; Surgeon, Royal Victoria Eye and Ear Hospital; Professor of Ophthalmology, University College, Dublin; and Examiner in Ophthalmology, Dublin University.

"We can whole-heartedly recommend the Book to Students. The Coloured illustrations drawn by the Author are a valuable feature."—"Lancet."

Fifth Edition (Reprinted). With 29 Plates (mostly in colour) comprising 33 figures. Demy 8vo. 7s. 6d. net; post free 8s.

## LANDMARKS & SURFACE MARKINGS of the HUMAN BODY

By L. BATHE RAWLING, M.B., B.C., Cantab. F.R.C.S., Eng., Surgeon, Demonstrator of Operative Surgery, St. Bartholomew's Hospital; Examiner in Surgery to the University of Cambridge, etc.

"We can confidently recommend it to everyone as a handbook both for study and for reference."—"Edinburgh Medical Journal."

SIXTH EDITION. Thoroughly Revised. With 2 Plates and 89 other Illustrations. Demy 8vo. 22s. 6d. net; post free, 23s. 6d.

## HYGIENE AND PUBLIC HEALTH

By LOUIS C. PARKES, M.D., D.P.H., Lond. Univ., Consulting Sanitary Adviser to H.M. Office of Works, etc.; and HENRY R. KENWOOD, C.M.G., M.B., Edin., D.P.H., F.C.S., Chadwick Professor of Hygiene in the University of London, etc.

[Lewis's Practical Series.]

"One of the most readable books on this now large subject that one can put one's hand on . . . has been thoroughly revised."—"The Medical Officer."

FOURTH EDITION. Revised. With 11 Plates (6 Coloured) and 74 other Illustrations. Demy 8vo. 22s. 6d. net; postage 1s.

## MIND AND ITS DISORDERS

By W. H. B. STODDART, M.D., Lond., F.R.C.P., Lecturer on Mental Diseases, St. Thomas's Hospital; Examiner in Psychology and Mental Diseases to the University of London, etc.

" . . . still remains as a standard work on psychiatry by an asylum physician of wide experience."—"Lancet."

FOURTH EDITION. Thoroughly Revised and greatly Enlarged. With 74 Plates and many other new illustrations. Demy 8vo. 28s. net; post free, 29s. 3d.

## DISEASES OF THE NOSE AND THROAT.

By HERBERT TILLEY, B.S.Lond., F.R.C.S.Eng., Surgeon to the Ear and Throat Department, University College Hospital; Teacher of Laryngology and Otology, University of London.

"We congratulate the author upon the success of his arduous task of revision, and have no hesitation in recommending his book to the practitioner as well as to the prospective specialist."—"Lancet."

[Lewis's Practical Series.]

ELEVENTH EDITION. Thoroughly Revised. Crown 8vo. 16s. net; postage 9d.

## ELEMENTS OF PRACTICAL MEDICINE.

By ALFRED H. CARTER, M.D., M.Sc., F.R.C.P.Lond., Formerly Professor of Medicine, University of Birmingham. Revised and edited by A. G. Gibson, M.D.Oxon, M.A., F.R.C.P., Lecturer in Morbid Anatomy, University of Oxford, etc.

"It provides a safe guide to Clinical Medicine, and possesses the merit of simplicity. . . . The book is well deserving of the wide circulation that it has obtained."—"The Lancet."

SIXTH EDITION. Revised and Enlarged. With Illustrations.

Demy 8vo. 21s. net; post free 22s.

## ANÆSTHETICS: Their Uses and Administration.

By DUDLEY W. BUXTON, M.D., B.S., M.R.C.P., Consulting Anæsthetist and late Lecturer to University College Hospital; Consulting Anæsthetist to the National Hospital for Paralysis and Epilepsy, Queen Square, Senior Anæsthetist and Lecturer on Anæsthetics in the Royal Dental Hospital of London, etc.

"The descriptions of apparatus and the illustrations in the book are extremely good. . . . The book is one to be read with profit and pleasure."—"Lancet."

[Lewis's Practical Series.]

\*.\* Complete Catalogue post free on application.

LONDON: H K LEWIS & CO., Ltd., 136 Cower St. & 24 Cower Place, W C.1

Telegrams—"Publicavit, Eusroad, London."

Telephone—Museum 1072.

# E. & S. LIVINGSTONE, MEDICAL PUBLISHERS

16 & 17 TEVIOT PLACE, EDINBURGH.

In Preparation for Early Publication.

## KERR, FERGUSON, HENDRY AND YOUNG. COMBINED TEXT-BOOK OF OBSTETRICS AND GYNECOLOGY.

Royal 8vo. 864 pp., 400 Illustrations. 35s. net.  
*Full Particulars on Application.*

## CHIENE'S HANDBOOK OF SURGERY. (Companion Volume to Jack's Medicine).

Crown 8vo. 600 pp. Fully Illustrated. 12s. 6d. net.  
*Full particulars sent post free on application.*

### Selected Publications.

#### SPECIALITIES AND TEXT-BOOKS.

**Bianchi—The Mechanism of the Brain and the Function of the Frontal Lobes.**  
Translated by McDonald. Royal 8vo. 368 pp.  
60 Illustrations. 25s. net.

**Glaister—Medical Jurisprudence and Toxicology.** 4th Edition. Demy 8vo. 918 pp.,  
158 Illustrations. 30s. net

**Robertson—Therapeutic Immunization.**  
Demy 8vo. 286 pp. 15s. net.

**Cushman—Chemistry and Civilisation.**  
Demy 8vo. 151 pp., 15s. net.

**Kraepelin—Dementia Praecox and Paraphrenia.** Trans. by Barclay. Edited by Robertson. Demy 8vo., 340 pp. 43 Illustrations. 17s 6d. net.

ALSO

**Manic Depressive Insanity and Paranoia.**  
Demy 8vo., 295 pp. 49 Illus. 21s. net.

**Maynard—Manual of Ophthalmic Practice.**  
Royal 8vo., 337 pp. 191 Illus. 25s. net.

ALSO

**Ophthalmic Operations.**  
2nd Edition. 267 pp. 142 Illus. 21s. net.

**Sidis—Symptomatology, Psychognosis, and Diagnosis of Psychopathic Diseases.**  
Demy 8vo., 467 pp. 21s. net.

**Talbot—Developmental Pathology.**  
Med. 8vo., 451 pp. 346 Illus. 25s. net.

#### HANDBOOKS.

**Burnet—Diseases of Children.**  
2nd Edition. 424 pp. Illus. 10s 6d. net.

**Gardiner—Skin Diseases.**  
180 pp. 46 Illus. 6s. 6d. net.

**Jack—Handbook of Medicine.**  
6th Edition. 568 pp. 28 Illus. 12s. 6d. net.

**Ross—Anæsthetics.**  
226 pp. 54 Illus. 3s. net.

**Syme—Nose, Throat and Ear.**  
337 pp. 26 Illus. 9s. 6d. net.

**Whittaker, J. R.—Anatomy of the Brain and Spinal Cord.** 5th Edition. 278 pp. 103 Illus. 12s. 6d. net.

**Whittaker, C. R.—Surgical Anatomy.**  
3rd Edition. 440 pp. 90 Illus. 15s. net.

ALSO

**Nerves of the Human Body.**  
Demy 8vo. Illustrated with Plates. 5s. net.

## BALLANTYNE'S POCKET BOOK OF OPHTHALMOLOGY.

120 pp. of type. Interleaved. 5s. 6d. net.

**Hewat—Examination of the Urine and other Clinical Side-room Methods.** 6th Edition. Illustrated. 2s. 6d. net.

**MacDonald—Pocket Prescriber, etc.**  
7th Edition. 200 pp. 3s. net.

## CATHCART—TRAINING AND HOW TO KEEP ALWAYS FIT.

52 pp. Attractive Cover. 2s. net.

FULL CATALOGUE SENT POST FREE ON APPLICATION.

E. & S. LIVINGSTONE, 16 and 17 TEVIOT PLACE, EDINBURGH.



## CONTENTS.

	Page
ORIGINAL COMMUNICATIONS—	
Fibrocystic Disease of the Tibia. By WILLIAM PEARSON	341
Deep Roentgen Therapy. By WILLIAM DOOLIN	... 357
MEDICAL EDUCATION	... 362
REVIEWS	... 371
ABSTRACTS OF CURRENT LITERATURE	... 382
PERIODICALS RECEIVED	... 387

## St. Andrew's Hospital

**FOR MENTAL DISEASES,  
NORTHAMPTON.**

of both Sexes. The Hospital, its branches (including a Seaside Home at Llanfairfechan, North Wales), and its numerous villas are surrounded by nearly a thousand acres of Park and Farm. Voluntary Boarders without Certificates received.

For particulars apply to DANIEL F. RAMBAUT, M.A., M.D., the Medical Superintendent.

President—The Right Hon. the  
EARL SPENCER, K.G.

This Registered Hospital receives  
for treatment PRIVATE PATIENTS  
of the UPPER and MIDDLE CLASSES

## HEIGHAM HALL, NORWICH.

Telephone: FOR UPPER AND MIDDLE CLASSES. 80 NORWICH.

A Private Home for Cure of Ladies and Gentlemen suffering from Nervous and Mental Diseases. Extensive pleasure grounds. Private Suites of Rooms with Special Attendants available. Boarders received without certificates.

Terms from 4 guineas weekly. Patients sent for.

Apply, DR. G. STEVENS POPE or MRS. POPE, Resident Licensees.

TRADE MARK 'SOLOID' BRAND  
ANTISEPTICS



SPEEDY — ACCURATE — CONVENIENT

Simply need to be dissolved in a definite quantity of water.

Portable, convenient, and exceptional in accuracy and purity.


Their unique shape, and the distinctive colouring of those containing poisons, prevent confusion with products intended for internal administration.

'SOLOID' products are prepared in the approved strengths of all the regularly used agents.

Example:— 'Soloid' Corrosive Sublimate, gr. 8.75. 1 in 1 pint of water = 1 in 1000 solution.



BURROUGHS WELLCOME & CO.  
LONDON



# THE IRISH JOURNAL OF MEDICAL SCIENCE

FORMERLY  
THE DUBLIN JOURNAL OF MEDICAL SCIENCE.

THE OFFICIAL ORGAN OF THE ROYAL ACADEMY  
OF MEDICINE IN IRELAND

---


---

FIFTH SERIES.      No. 8.      OCTOBER, 1922

---

---

*Original Communications.*



## A CASE OF FIBROCYSTIC DISEASE OF ~~THE~~ TIBIA; TREATED BY RESECTION AND BONE-GRAFTING.\*

BY WILLIAM PEARSON.

OSTEITIS Fibrosa or Fibrocystic disease of bone is a rare disease, but one of much interest to pathologists and surgeons. The condition receives scant notice in surgical text books, and its etiology is obscure. It has, however, been thoroughly reviewed in excellent papers by Bloodgood<sup>1</sup> and Elmslie<sup>2</sup>.

From a study of 89 cases classified as bone cysts Bloodgood distinguished two main groups :

- (a) True bone cysts, which have a definite relation to osteitis fibrosa, and
- (b) Cysts in the medullary cavity due to other conditions.

The first group contained 69 cases, which he divides into six varieties :

1. Single cyst, bony shell, no connective-tissue lining.

Read in the Section of Surgery, Royal Academy of Medicine in Ireland, April 28th, 1922.

2. Single cyst, with definite connective-tissue lining.
3. A small cyst or cysts in a solid mass of osteitis fibrosa.
4. No cysts, the bone shell being filled with a solid mass of osteitis fibrosa.
5. Multilocular cysts.
6. Miscellaneous.

"These cases," he says, "have in common the presence of an inflammatory tissue in the medullary cavity (osteitis fibrosa), which, with or without cyst formation, is replacing the marrow and cancellous bone, producing absorption of the outer table, and associated with more or less distension of the thin bone shell. Unless there has been a pathological fracture, or the cysts are of huge size and long duration, the periosteum is normal and there is little or no new bone formation." "In a few of these cases, here and there, islands of cartilage are found . . . . . but cartilage is never present in sufficient quantity to justify the conclusion that the cyst is due to the liquefaction of a primary or original solid area of cartilage."

"The fluid content of the cyst is never distinctly hæmorrhagic: it is usually thin, dark brown in colour, and contains microscopically blood pigment and blood cells." Giant-cells are often found near the lamellae of the bone shell, but except in one or two cases, have never been found in sufficient numbers to characterise the picture as a giant-cell sarcoma. They are to be regarded as giant-cell osteoclasts producing bone absorption.

The condition may arise in many situations, but occurs most often in the upper ends of the humerus and femur, and in the tibia.

All observers agree that it is essentially a disease of youth. It usually starts before the age of 15, but is so insidious in its onset and progress, that it may not be recognised until a much later period. Only five of the 69 cases reviewed by Bloodgood were over 20 years of age when the disease was first noted.

The clinical manifestations in their order of frequency are pathological fracture, pain and swelling. Swelling will be more frequently noted when the disease affects a superficial bone such as the tibia. Disability of the limb may be noted,





Fig. 1.




Fig. 2.



Fig. 3.



Fig 1.



but usually only after an injury or a definite fracture. The condition cannot be differentiated without the aid of skiagrams.

Theories differ widely in regard to the etiology: Virchow<sup>3</sup> considered that the cysts developed from enchondromata by degeneration.

Bloodgood and other observers regard the condition as an inflammatory one, the medullary tissue being replaced by connective tissue with or without cyst formation. Consequently, he suggests the terms chronic osteomyelitis cystica or solida as appropriate.

Elmslie discusses the matter at length, and considers that in the present state of our knowledge, we cannot decide whether fibrocystic conditions in bone constitute one disease or several. He rejects Virchow's theory, and does not agree that the condition is inflammatory. He suggests as a working hypothesis "that there is a general disturbance in the processes of growth or alteration in the bone. "That in the course of this there is (1) a fibrous metaplasia of the marrow; (2) an excessive formation and activity of the giant-cells; (3) an active absorption of bone, with coincident formation of new bone and cartilage. That in places the fibrous metaplasia may be most evident leading to the formation of fibrous masses; in other places cysts may be produced, either by the action of giant cells, or by degeneration in the fibrous tissue; and that in still other places there may be abundant formation of bone—which may or may not be calcified—and of cartilage or fibrocartilage. . . . ."

The case recorded here is that of S.A., male, aged 20 years, who came under my care suffering from a swelling of the left leg which had been regarded as a sarcoma of the tibia. The swelling had been first noticed four years previously, it had slowly increased in size, and had only recently become slightly painful. Examination showed a fairly smooth fusiform swelling in the middle third of the shaft of the tibia, with some anterior bowing of the bone. The swelling was slightly tender on firm pressure, and did not involve any of the soft tissues which were freely movable over it. The skin was normal and the function of the limb was not impaired.

The clinical symptoms and features of the case rendered

the diagnosis of sarcoma uncertain, and skiagrams showed the condition to be typical of fibrocystic disease (see Fig. 1).

A subperiosteal resection of the shaft of the tibia was performed, the bone being divided above and below the limits of the disease, and somewhat more than the middle third of the shaft was removed.

In order to determine what amount of regeneration would follow subperiosteal resection in the case of a young adult in the absence of sepsis no bone graft was inserted at this stage, and the wound was closed without drainage. Skiagrams taken one and two months after resection (see Figs. 2 and 3) showed that practically no new bone had formed, save in the neighbourhood of the upper segment from which a slender cone of ossification extended downwards for a distance of one inch. Elsewhere only very light flakes of bone were seen scattered here and there in the periosteal tube. It is interesting to note that these flakes all lay in the long axis of the periosteum.

Three months after resection a graft was taken from the crest of the right tibia and inserted as an intramedullary peg, being embedded inside the periosteal tube between the ends of the tibia. This tube had become thickened almost to the size of a normal tibia and on section was firm, smooth and homogeneous, with very little vascularity, giving one the impression of soft cartilage.

Nine months later the patient was able to walk well without support, and no deformity or want of symmetry could be detected.

The progress of the case may be seen by a study of the radiograms taken at intervals after operation. They demonstrate that although the graft was successful, and became firmly united to the tibia at each end, it shows little increase in size at the end of nine months, and that regeneration of a new tibial shaft has been effected chiefly by ossification which proceeded throughout the substance of the thickened periosteal tube in which the graft was embedded (see Figs. 4, 5, 6, 7, 8 and 9). This variation from the usual behaviour of a successful "bridge" graft is, I think, due to the fact that the periosteum between the ends of the tibia was intact




Fig. 5.



Fig. 6.





Fig. 7.



Fig. 8.

in this case, but this does not mean that the new bone was developed directly from the periosteum.

The precise mode of action of the graft is open to question. A close study of the skiagrams suggests that it has been both osteogenetic and osteoconductive, chiefly the latter—and that the osteoblasts for the most part have been furnished by emigration from the ends of the tibia by way of the graft.

Examination of the gross specimen on section showed that the medulla and normal cancellous tissue were replaced by a spongy-looking mottled red and yellow substance, in which small translucent areas were present indicative of early cyst formation. Absorption of the cortex, and expansion and bowing of the shaft were also very evident. Microscopically the tissue consisted of a broken network of bone lamellæ, the spaces between which were filled with a fibrous connective tissue (osteitis fibrosa) in which minute cystic cavities were present in places. Here and there giant-cell osteoclasts were present, producing absorption of bone on the periphery of the lamellæ. The microscopic appearances were identical with those illustrated in the photo-micrograph which is reproduced from Bloodgood's communication.

#### DESCRIPTION OF PLATES.

Fig. 1. Osteitis fibrosa of tibia. The middle third of the shaft is expanded and presents a finely mottled appearance indicating irregular bone absorption, and suggesting the development of numerous small cavities or cysts in its substance. The cortex is greatly thinned along the anterior surface where the expansion is slightly irregular, but at no points has the disease broken through or invaded the periosteum. The disease involves a much greater extent of the anterior than of the posterior aspect of the bone. At either end it is limited by a zone of condensation which separates it from the normal medullary cavity; this is more marked at the lower end. The shaft shows a considerable degree of antero-posterior bowing.

Fig. 2. Tibia one month after subperiosteal resection of shaft. Very faint evidence of callus formation in connection with the upper segment.

Fig. 3. Two months after resection. A cone of callus is seen projecting from the upper end. Slight evidences of callus formation in the form of longitudinally arranged flakes of bone are seen scattered throughout the periosteal tube.

Fig. 4. Three weeks after grafting. The graft is united at each end to the tibia, but no trace of further bone formation in the periosteal tube.

Fig. 5. Eight weeks after grafting. The graft is practically unchanged, but there are signs of ossification beginning throughout the periosteal tube. The cone of ossification proceeding from the upper tibial segment has increased in length and thickness and shows longitudinal striation. A faint zone of rarefaction is seen round the lower end of the graft which suggests that it is becoming loose at this point, but subsequent skiagrams show that this is not so.

Fig. 6. Four months after grafting. The graft is firmly united to the tibia. The sharp edges of the graft are becoming smoothed off, and the extremities are losing definition and becoming merged in the structure of the tibia. Ossification is proceeding throughout the periosteal tube surrounding the graft; this ossification is not quite uniform near the upper end, where a few lighter areas are seen in the callus.

Fig. 7. Five months after grafting. Similar to Fig. 6, with ossification slightly more advanced. Daily massage had been employed for four weeks.

Fig. 8. Eight months after grafting. During the previous three months the leg had been encased in plaster of Paris and patient had been using crutches. Very little change noticeable.

Fig. 9. Nine months after grafting. During the previous month patient had been walking on the limb and receiving daily massage. The "periosteal" callus has increased in density and definition so that the normal outline and contour of the tibia have been restored. The differentiation between graft and callus is still quite distinct and a line of "demarcation" (Fig. 9) is seen between the two in places. This is due to the fact that the graft was covered with its own periosteum on this aspect, so that a fibrous membrane intervened between the bone of the graft and the ossifying periosteal tube. This would appear to indicate that grafts placed intraperiosteally should preferably be deprived of their own periosteal covering.

#### *References :*

1. Bloodgood : *Annals of Surgery*, August, 1910.
2. Elmslie : *British Journal of Surgery*, July, 1914.
3. Virchow : *Virchow's Archives*, LXX, 502






Fig. 9.



Fig. 10.

Photomicrograph of Osteitis Fibrosa (Bloodgood). The wall of the cyst is lined by a well-marked layer of cellular fibrous connective-tissue, which is replacing the bony framework. Giant-cell osteoclasts may be seen in places producing bone absorption.


# SURGICAL INSTRUMENTS.



63659.

## MIDWIFERY BAG.

In Black or Brown Cowhide.  
Lining removable.  
16 in. ... £4 4 0



C3650.

## SURGICAL OR MIDWIFERY BAG.


In Black or Brown Cowhide.  
Lining removable.  
12 in. ... £2 16 0  
14 in. ... £3 3 0  
16 in. ... £3 12 0



C3960.

## VAGINAL SPECULA.


5/- each.  
Set of 4, 18/6.



C3652.

## MIDWIFERY BAG.

Black or Brown Cowhide.  
Lining removable.  
12 in., 30/-; 15 in., 32/-; 16 in., 34/-.



C3651.

## MIDWIFERY OR SURGICAL CASE.

Black or Brown Cowhide.  
17 x 10 x 5 in. ... £2 16 0  
Fittings Extra.




C3955.

## AUWARD'S SPECULA.

18/-.


**REDUCED PRICES.**—Our 800 page Illustrated Catalogue, giving present-day prices, will be sent free on application.



C3668.

## NICKEL-PLATED COPPER STERILIZER.

Complete as illustrated.  
16 x 4 x 3½ ins. ... £2 2 0.



C3875.

## SET OF DOUBLE-ENDED DILATORS, 35/-, HEGAR'S.



C3954.

## VAGINAL SPECULA. SIMS.


3 sizes,  
5/6 each.



C3855.

## FLUSHING CURETTE, RHEINSTATER'S, 7/6.

**The SURGICAL MANUFACTURING CO., Ltd.**  
83-85 MORTIMER STREET, LONDON, W. 1.



System II.

## DOMEN BELTS

For All Abdominal Troubles

Speaking of System II.—the belt illustrated—the Medical Annual 1917 says:—"It is in Enteroptosis that we have used it most frequently and with invariably good results."

Domen Belts Co., Ltd.  
456 Strand, London, W.C.2.

## HORLICK'S MALTED MILK

THE ORIGINAL.

THE IDEAL ALL-WEATHER  
FOOD-DRINK.

Made in England.

Horlick's Malted Milk is meat and drink in one, because it contains all necessary nutritive elements in the correct proportions demanded by Nature to support life and maintain health. Composed of the extracts of selected malted barley and wheat flour, combined with pure, Pasteurised cow's milk, it is perfectly digestible and rapidly assimilated, while Vitamines are supplied by both the milk and the grain.

READY IN A MOMENT BY STIRRING BRISKLY  
IN HOT OR COLD WATER ONLY.

Always specify "HORLICK'S"

Free Samples sent to Members of the  
Profession on application to

HORLICK'S MALTED MILK CO.,  
Slough, Bucks., England.

For Use in Bath and Toilet Basin.

## SULPHAQUA

NASCENT SULPHUR  
CHARGES

Largely prescribed in

GOUT, RHEUMATISM, ECZEMA, THE S.P. CHARGES CO.  
SCABIES, and all SKIN DISEASES. St Helens, Lancs

Baths prepared with SULPHAQUA possess powerful antiseptic, antiparasitic, and analgic properties. They relieve intense itching and pain, are without objectionable odour, and do not blacken the paint of domestic baths.

SULPHAQUA SOAP.—Extremely useful in the treatment of Acne and Seborrhœa of the Scalp and Eczematous and other skin troubles. In boxes of ½ and 1 doz. Bath Charges, 2 doz. Toilet Charges, and ¼ doz. Soap Tablets. Advertised only to the Profession.

The STANDARD  
HÆMATINIC.

"The employment of Bi-Palatinoids places the administration of exceedingly unstable compounds like ferrous carbonate, phosphate and arsenate, upon a sounder and more scientific basis."—The Lancet.

## "BI-PALATINOIDS"

(No. 500)

OF NASCENT FERROUS CARBONATE.

Sample and Literature  
on application.

OPPENHEIMER, SON & CO., LTD.,  
179 Queen Victoria Street, London, E.C. 4.  
Entirely British House since Foundation in 1891

# AN IMPRESSION OF "DEEP ROENTGEN THERAPY" IN SURGICAL WORK.

(FRANKFURT-AM-MAIN).

BY WILLIAM DOOLIN.

THE deep therapy apparatus for surgical purposes is situated in one wing of the surgical klinik (350 beds), under the direction of Professor Schmieden, who has been in control here since 1919. The Röntgen specialist in charge is Dr. Hohlfelder, working with the following staff, who are employed exclusively in the *x-ray* treatment of the surgical patients: three doctors (whole time assistants), four lady assistants, and six "Röntgenswestern" (*x-ray* sisters); each of these must spend not less than a six months' apprenticeship in the *x-ray* department, and may not work in any other department.

The *x-ray* department consists of one complete wing, of which the treatment rooms cover an area of about one-third the floor space. Four, or at most, six patients can be treated at a time. The *x-ray* diagnostic division, situated in the same wing, has an entirely different apparatus and personnel, under Dr. Hohlfelder's supervision. Plates and diagnosis reports are kept in the Bibliothek, or library.

It will be clearest if we follow the course of a patient who has been referred to the klinik as a suspected case of cancer of the stomach.

Such a patient passes through the following examination:—

1. The usual personal examination (physical and history taking) is carried out by at least two members of the surgical staff.
2. Differential diagnostic test meals, examination of fæces for occult blood, etc., are carried out by the resident pathologist.
3. A complete diagnostic Barium-meal *x-ray* examination is carried out by Dr. Hohlfelder: and a routine *x-ray* examination of thorax and bony skeleton is made for possible metastases.



4. A complete blood examination is made, and tabulated.
5. Patient is weighed, stripped.

When the results of these separate examinations have been scheduled into one "protocol," or case-history, the patient is submitted to the Professor, who, on the evidence before him, decides as to whether the patient is operable or not. The patient may be put at once into one of three classes of operability. 1.—Operable. 2.—Doubtful. 3.—Hopeless. If operable, he is admitted to hospital for operation and post operative *x-ray* treatment.

If in-operable, he is advised to stop in lodgings in the town, and to attend as an extern, for *x-ray* treatment.

Thus, all patients receive *x-ray* treatment, no matter to what class of operability they belong.

Treatment is carried out as follows :—

The patient comes fasting ; he receives morphin-scopolamin and then lies on the couch under continuous *x-ray* exposure for eight *x-ray* hours. (This may mean eight hours by the clock, with one tube, or four hours with two tubes.) The tumour area is irradiated for half the treatment period through the anterior abdominal wall. Then the patient is turned over on his face, and the tumour area is attacked from behind for the remaining four hours. The cancerous tissue thus receives a uniform irradiation of 2 m.a. per minute for eight hours. They calculate with an astonishing degree of mathematical accuracy the depths to which each individual ray will penetrate the tumour : the aim of the treatment is to expose each scrap of the tumour to a uniform irradiation during the eight hours. The *x-ray* dosage is estimated and controlled throughout by the Iontoquantimeter.

At the conclusion of the treatment, a new blood examination is carried out and tabulated. For the next two or three days the patient is often very sick and depressed : this reaction is due to two factors : (1) the massive *x-ray* dose has a markedly deteriorating influence on the circulating bloodcells, from which they take some three weeks to recover, and (2) into this weakened circulation are rapidly absorbed the toxic products of the necrosed cancer-mass.

The patient returns on the eighth day for re-examination,



when he is again examined by the two assistants who first saw him, is again weighed, a barium-meal examination is repeated, and any shrinkage of the growth is noted; the blood picture is again taken and compared, chemical meal tests and blood faeces examination repeated. If the patient had been originally put in the "doubtful" class, a further consultation is now held with the Professor as to whether an operation at this stage is likely to be of benefit, or not. He then goes home, to return again three weeks later, when he is re-examined in the same detail, and when it is decided whether he is to have a second course of *x-ray* treatment or not.

Patients who have been operated on may receive after operation two *x-ray* exposures, with one month's interval between each. Patients, who are hopeless from the operative point of view, with widely spread metastases, receive exposures depending on the extent and degree of metastases present.

Although a specific case of gastric cancer has been cited as an example, I would like to point out that this method of examination, control, and consultation, applies to all cases of surgical cancer in the klinik, no matter in what organ the tumour be situated. It so happens that, owing to Professor Schmieden's individual reputation, one sees more cases of abdominal and rectal cancer in his klinik than of any other organ.

Uterine cases, are of course, never to be seen in the surgical klinik. They are in the gynecological klinik, a separate building, under the control of Professor Seitz, who has just been called from Erlangen to Frankfurt, and who has his own separate *x-ray* installation and "team" for his uterine cancer cases.

#### *General Observations on Cancer Treatment.*

All cancer patients, whenever possible, are first submitted to operation, and then to *x-ray*. Pre-operative *x-ray* treatment is frequently utilised. Professor Schmieden greatly favours this "combined method" of treatment.

Sarcomata are treated exclusively by *x-ray*.

Surgical tuberculosis cases are very rarely operated on; but are treated by *x-ray*, by quartz-lamp, by insolation, etc., and are kept on an open air verandah, day and night.

With the utmost honesty, our instructors told us they could give us no definite pronouncement as to the precise present value of their cancer therapy. The method is in use in this klinik only since 1917, and four years' experience does not constitute a sufficient period of time in which to speak with authority of end results. Their attitude to the problem is thus expressed :—

In Germany, with these newer methods, we are certainly getting better cancer statistics than with our pre-war surgery, but there is still immense room for improvement. This improvement will come gradually with the use of the intensive anti-cancer campaign now being carried out by means of articles in the lay press, and of lectures to the lay public; our general practitioners are sending us cases earlier now than formerly, and we are further educating ourselves as we become better acquainted with our *x-ray* methods. This last means is still in its infancy, and we have much work yet to do before we bring it to perfection. There is as yet no specific cure for cancer, in the Salvarsan sense.

When we had explained to Professor Schmieden the different working system of Irish and German hospitals, he gave us as his distinct opinion that the use of the "deep *x-ray* treatment" for a small hospital was not practicable. "To get good results economically," he emphasised, "the method must be applied on the grand scale, and with a picked staff of whole-time technical experts. In Dublin, one "symmetrie" or "I.R.A." apparatus in a central clinical institute with a properly organised personnel, should suffice for 300-400 surgical beds. To use the apparatus in a small hospital means prohibitive expense. It takes at least six months' study for each individual member of the working team to become sufficiently familiar with the method to obtain satisfactory results.

Our enquiries showed us that even in Germany, the method is still *sub judice*, to this extent, that while all their authorities are at present satisfied that the combination of operative and *x-ray* treatment is giving better cancer results than those heretofore obtained, there still remains a widely expressed difference of opinion as to the actual methods of technique and apparatus to be employed. In other words, there is at

present no standard technique, and no standard apparatus. In Schmieden's klinik (as also in that of Professor Frankl, at Vienna, who was recently in Dublin), the "Symmetrie" apparatus is the machine in use. In Erlangen, Professor Wentz uses an "I.R.A." apparatus, whilst in Heidelberg, Enderlen employs yet another. All these workers are working on the same basic principles, each trying out some modification of technique of his own, in the endeavour to arrive at a definite maximum standard of technique, regardless of cost, which is borne by the state.

---

## MEDICAL EDUCATION.

THE system of medical teaching in Ireland differs from that in England in important particulars. In London each clinical hospital has its attached medical school, fully equipped, which educates the students of that hospital and very seldom those of any other. In Dublin, on the contrary, the hospitals and schools are entirely separate, and a student of any school is free to enter for the whole or any part of his course at any school or hospital he pleases.

### *Cost of Education in Ireland.*

School of Physic, Dublin University	...	£124	17
Royal College of Surgeons' School	...	124	19
National University Colleges	...	124	19
Queen's University, Belfast	...	105	0

### *Cost of Diplomas or Degrees.*

Dublin University—£27 (to this must be added £83 4s., the cost of obtaining an Arts degree).			
National University	...	£19	0
Queen's University, Belfast	...	19	19
Conjoint Royal Colleges	...	42	0
Apothecaries' Hall	...	22	1

Thus, the absolute payment will amount to somewhere between £125 and £233 1s., according to the course chosen. For the Conjoint Colleges the entire cost is £166 19s., taking the minimum mode of payment. So that, assuming that extras or voluntary costs are incurred, the total will vary, say, from £170 to £200.

All the schools require fees for each course to be paid in advance.

Women are admitted to all the courses, degrees and licences on the same terms as men.

*Date of Entry*

The entry of names and commencement of study in Ireland is *supposed* to date from the 1st of October in each year, but entries are accepted for some weeks later. The student must attend three-fourths of the lectures delivered.

*Preliminary Examinations.*

The first act of the student is to pass a preliminary examination. The next is to commence medical study. This he does by entering for lectures at a medical school. From the school registrar he gets a form of certificate, and his third act is to take it or send it to the Branch Medical Council, 35 Dawson Street, Dublin, unless, as is usually the case, this duty is undertaken for him by the school registrar. He is thereupon placed upon the Register of Medical Students (without fee), and his period of study counts from that date.

The only preliminary examination held specially for medical students is that held conjointly by the Royal Colleges of Physicians and Surgeons, but other examinations—e.g., the public entrance at Trinity College, the matriculation of the National and Queen's Universities, and the Intermediate Examination passes in the required subjects, are accepted as equivalent.

The subjects of examination as prescribed by the General Medical Council are as follow :—(1) English language, including a specified author ; dictation, grammar, and composition ; also parsing and analysis from a book specified ; (2) Latin, including grammar, translation from specified authors, and translation of easy passages not taken from such authors. (3) Elements of mathematics, comprising (a) arithmetic, including vulgar and decimal fractions ; (b) algebra, including simple equations ; (c) geometry, Euclid, Books I., II., III., with easy deductions. (4) One of the following optional subjects :—(a) Greek ; (b) French ; (c) German.

## THE UNIVERSITIES.

*The University of Dublin.*

The University of Dublin grant the degrees of M.B., B.Ch., and B.A.O. to students who have obtained their

B.A. degree, and who have been for at least five academic years on the books of the Medical School, and the higher degrees of M.D., M.Ch., and M.A.O., to graduates of certain standing who hold the degrees of M.B., B.Ch., and B.A.O. It does not grant degrees to any but graduates in Arts, and consequently its degrees hold a high rank and are sought for by those who look forward to occupying the best positions in the profession. Diplomas, entitling to registration, are given on certain conditions to non-graduates.

The expense of obtaining the degrees of M.B., B.Ch., and B.A.O. is approximately as follows:—Lectures, £73 10s.; Hospitals, £55 13s.; Degree Fees, £17—total, £146 3s.

The expense of the B.A. degree, amounting altogether to £101, should be added, making the total cost £247 3s.

In addition to its ordinary qualifications the University grants the following degrees and diplomas:—

*Doctor of Medicine.*—To obtain this the candidate must have passed the final examinations and be of M.A. standing. He must send in a thesis for approval. Subsequently the Regius Professor of Physic and an assessor will discuss with him questions connected with the thesis and may examine him *viva voce* on medical subjects of a more general nature. Fee for this degree, £20.

*Master of Surgery.*—The candidate must be a Bachelor in Surgery of three years' standing, and must then pass an examination in clinical surgery, operative surgery, surgical pathology, surgery, and surgical anatomy (on the dead subject). Fee for this degree, £11.

*Master in Obstetric Science.*—The candidate must be a B.A.O. of two years' standing, and must produce satisfactory evidence of having been engaged in the study of obstetrics for two years. He is then required to pass an examination in the following subjects:—Practice of midwifery, gynaecology, anatomy of female pelvis, and elementary embryology and clinical gynaecology. Fee for this degree, £15.

*Diploma in Gynaecology and Obstetrics.*—The candidate need not be a graduate in Arts, but must have been a registered practitioner for at least twelve months and have spent at least one year after registration in the study of obstetrics and gynaecology. During this year's course he must have



been resident for six months in Trinity College and for six months in the Rotunda Hospital.

*Diploma in Public Health.*—The candidate must be a registered medical practitioner; must have completed, subsequent to obtaining a registrable qualification, four months' practical instruction in a laboratory in practical work in chemistry and bacteriology applied to public health; he must have studied, practically, outdoor sanitary work for six months under an approved officer of health; and have attended, after qualification, for three months the practice of a hospital for infectious diseases.

*Degree in Dental Science.*—Candidates for the B.Dent.Sc. degree must have taken a degree of Arts and must have had their names in the books of the Medical School for four years. Three examinations must be passed—namely, the Preliminary Scientific at the end of the first year; the Intermediate at the end of the third year; and the Final Dental at the end of the fourth year. The total fees are £309 4s. Candidates for the degree of Master in Dental Science must be Bachelors in Dental Science of at least one year's standing. They will be required to pass an examination in Pathology and Bacteriology, and either to carry out Dental work of an advanced character to the satisfaction of the examiners, or to present a thesis, to be approved of by them, giving evidence of original research on some subject connected with dentistry.

The fee for the M.Dent.Sc. examination is £5, and the fee for the degree is £10.

Full particulars regarding the Medical and Dental Courses and a prospectus of the Courses for the Diploma in Public Health may be obtained by application to the Registrar of the School of Physic, Trinity College, Dublin.

### *National University of Ireland.*

The National University of Ireland confers the degrees of M.B., B.Ch., and B.A.O. on students who have followed the prescribed course for five academic years, and passed the prescribed examinations. At least three years must be spent at one of the constituent colleges of the University—namely, the University Colleges of Dublin, Cork, and Gal-

way. The University also confers the degrees of M.D., M.Ch., M.A.O., B.Sc. (Public Health), D.Sc. (Public Health), B.D.S., Primary degrees, and M.D.S.

The expense of obtaining the degrees of M.B., B.Ch., and B.A.O., in University College, Dublin, is approximately—Lectures, £91; Hospitals, £54 14s.; University fees, £21; total, £169.

The conditions for the higher degrees are :—

*Doctor in Medicine.*—Candidates may present themselves for the examination for this degree after an interval of three academic years from the time of obtaining the M.B., B.Ch., B.A.O. degrees; but in the case of candidates who have obtained a degree of the University in the Faculty of Arts, or the Faculty of Science, an interval of two academical years is sufficient.

Candidates must at the same time produce a certificate of having been, for at least, two academical years, engaged in hospital or private medical, surgical, or obstetrical practice respectively, or in the military or naval medical service.

Candidates at this examination must answer in Medicine and Pathology.

*Master in Surgery.*—The following are the subjects of examination :—

Surgery, theoretical and practical, including Ophthalmology and Otology; Surgical Pathology; Surgical Anatomy and Operative Surgery, with the use of Surgical Instruments and Appliances. The other conditions are the same for the M.D. degree.

*Master in Obstetrics.*—Academic standing is as for the two previous degrees. Each candidate must furnish satisfactory evidence that since he has (1) had personal charge of at least *twenty* cases of labour; and (2) attended during a period of three months the practice of a clinical hospital for diseases of women where at least six beds are in constant occupation or in a special ward of a general hospital, where such cases only are treated, and containing at least six beds in constant occupation.

Candidates at this examination must answer in the following subjects: Midwifery, Diseases of Women and Children, Pathology, the Use of Instruments and Appliances. Fee

for each of the above degrees is £10 10s. In the case of graduates who have matriculated in the Royal University the fee is £5.

*Bachelor of Science, Public Health.*—A candidate shall not be admitted to receive the degree unless he—(a) shall have received the degrees of M.B., B.Ch., and B.A.O., at least one year previously; (b) shall have pursued an approved course of study in the Faculty of Medicine; and (c) shall have passed the prescribed examination. Fee, £7 10s.

*Courses.*—In addition to D.P.H. Course; (1) Special Pathology (three months); (2) Bacteriology second course (three months); (3) Advanced Course in Hygiene (three months).

*Doctor of Science, Public Health.*—The regulations are not yet published. Fee, £10 10s.

*Diploma in Public Health.*—This Diploma may be granted to matriculated or non-matriculated students of the University who shall have completed courses of study, approved for the purpose, and shall have passed the prescribed examinations; provided that it shall not be granted except to a registered medical practitioner.

The conditions and examinations are similar to those already quoted for the University of Dublin.

*Degrees in Dentistry.*—Candidates for the degree of B.D.S. shall be required to pass, after matriculation, four University Examinations—namely: A first University Examination; a Second University Examination as for Medical Students; a Third University Examination in Dental Surgery; a Final Examination for the degree of B.D.S. University Fees, £16 10s.

*Master of Dental Surgery.*—Candidates may present themselves for the examination for this degree after an interval of three academic years from the time of obtaining the B.D.S. degree. Fee, £10 10s.

The University also grants Diplomas in Mental Diseases and Tropical Medicine.

IRISH MEDICAL CORPORATIONS GRANTING  
DIPLOMAS.*Royal College of Physicians of Ireland.*

This College issues a Licence in Medicine and a Licence in Midwifery to registered medical practitioners.

*Licence in Medicine.*—The subjects of examination are—Practice of Medicine. Clinical Medicine, Pathology, Medical Jurisprudence, Midwifery, Hygiene and Therapeutics.

*Licence in Midwifery.*—The subjects of examination are—Gynæcology and Midwifery.

*Fees.*—Fee for the Licence to Practice Medicine, £15 15s. Fee for the Licence to Practise Midwifery, £5 5s.

*Membership.*—The Membership is open to University Graduates in Medicine and to Licentiates of the Royal Colleges of Physicians of the United Kingdom. The Examinations for Membership are held in February, June and November, and such other times as the President may appoint. Fee to Licentiates of the College, £21 ; to others, £36 15s. Special Examinations £10 10s. extra.

*Fellowship.*—The Fellowship is open to all Members of the College of one year's standing or over, irrespective of sex. Fee £60. Election is by ballot.

*Royal Colleges of Physicians and Surgeons.*

Examinations are held conjointly by the two Colleges. The course, as in other bodies, extends over five years, with examinations at the end of the first, second, third and final years. As in the English Colleges the subjects of the First Professional Examination may be studied either at a medical school or at an institution other than a medical school, which is recognised by the Colleges, after due inspection for instruction in these subjects. We recommend students to apply for the official programme to the Secretary of the Committee of Management, Royal College of Surgeons, or to the Registrar of either College. In the case of the Preliminary Examination seven clear days' notice must be given to the Secretary.

The Colleges also confer a Diploma in Public Health.

*Royal College of Surgeons in Ireland.*

This College grants a Licence in Surgery to registered medical practitioners.

The subjects, methods, times and places of examination are those of the surgical group of the Final Professional Examination of the Conjoint Board in Ireland of the Royal College of Physicians and the Royal College of Surgeons. Special examinations will not be granted under any circumstances.

The fee for examination for each admission is five guineas. The fee to be paid upon admission to the Licence in Surgery is twenty-five guineas.

Fees will not be returned under any circumstances.

*Fellowship.*—The examination for the fellowship is divided into two parts—namely, the Primary and the Final. The subjects of the primary examination are anatomy, including Dissections, Physiology and Histology. The subjects of the final examination are Surgery, including Surgical Anatomy and Pathology. Candidates must pass in all the subjects at one examination. The examinations are held three times in each year.

*Fees.*—The fees for examination are as follows :—(1) Primary examination, each admission, 5 guineas ; final examination, each admission, 5 guineas. (2) Of these examination fees ten guineas will be reckoned as part of the fee payable upon admission to the Fellowship. (3) The fee to be paid upon admission to the Fellowship is forty guineas, except when the candidate is a Licentiate of the College in which case the fee is twenty-five guineas.

*Licence in Dental Surgery.*—Candidates for the Licence in Dental Surgery are required to pass two professional examinations. They must have passed a recognised Preliminary Examination in general education, have been registered as medical or dental students by the General Medical Council, and have been engaged for two years in acquiring a practical familiarity with the details of mechanical dentistry under the instruction of a registered dentist, or under the direction of the superintendent of the mechanical department of a recognised dental hospital where the arrange-

ments for teaching mechanical dentistry are satisfactory to the Council of the College. The instruction may be commenced or attended before the candidates register as medical or dental students. One year's *bona-fide* apprenticeship with a registered dental practitioner, after being registered as a medical or dental student, may be counted as one of the four years of professional study required. There are special exemptions in the case of persons already holding a surgical or dental qualification.

*Apothecaries' Hall of Ireland.*

The Licence of this Hall is granted to students who present certificates of having completed the course of study as laid down in the curriculum and who pass the necessary examinations. The diplomas of the Apothecaries' Hall of Ireland entitle the holder to be registered as a practitioner in medicine, surgery, and midwifery, and possess the privileges of an apothecary.

There are three professional examinations, the total fees for which amount to 40 guineas. Women are eligible for the diploma.

Registered medical practitioners receive a diploma of the Hall upon passing an examination in the subject or subjects not covered by their previous qualification, and on paying a fee of £26 5s. (twenty-five guineas).

Each candidate before receiving his diploma must produce evidence that he has attained the age of 21.

Each candidate must produce evidence of having before entering on medical studies passed a preliminary examination in general education.

The details of the course of education required and syllabus of the examinations will be supplied on application to the Registrar at 40 Mary Street, Dublin.



# 'HYGEIA' NURSING HOME

## VENTNOR, ISLE OF WIGHT.



"Hygeia" Nursing Home is beautifully situated. Ventnor has probably the most equable climate in England. The Home is sheltered from the North and East by the downs and obtains the maximum amount of sunshine.

amount of sunshine, and is therefore particularly suitable for bed cases of "ante" or "post" sanatorium patients.

Every care is taken to see that diet is suited to the needs of each individual patient.

A Doctor is always at hand in cases of emergency, and trained nurses are on duty night and day.

Ventnor is 92 miles from London and is reached in about four hours.

For medical references, terms, etc., apply to the Secretary.

'Phone: 29 Ventnor.

Possessing advantages over any other preparation of sulphur available in therapeutics.

## Collosol Sulphur

(CROOKES)

Indicated in Acute, sub-acute and chronic Rheumatism, Rheumatoid Arthritis, Neuritis, Lumbago and Sciatica, Chronic skin affections attributable to auto-intoxation in which it acts as an intestinal antiseptic.

A 1% colloidal solution showing marked Brownian activity.

Collosol Sulphur readily combines with proteins in the intestinal tract and is entirely absorbed without the formation of sulphuretted hydrogen.

Clinical samples and literature to the profession on application.

### THE CROOKES LABORATORIES,

BRITISH COLLOIDS, LTD.,

22 Chenies Street, Tottenham Court Road, W.C.1

Telegrams: "Colosolly, Westcent, Lon."

Telephones: Museum, 3663-3697.

Eighth Edition. Fully Revised with additional Chapters.  
Over 360 Illustrations, many being new or re-drawn for this edition, and 11 Plates.  
21s. net. Postage 1s.

## **PYE'S SURGICAL HANDICRAFT.**

A Manual of Surgical Manipulations, Minor Surgery, and other Matters connected with the work of House Surgeons, Surgical Dressers, etc.

Edited and largely re-written by  
**W. H. CLAYTON-GREENE, C.B.E., B.A., M.B., B.Ch. Camb., F.R.C.S.**  
Surg. to St. Mary's Hosp.; Lecturer on Surgery in the Medical School, etc.

With special chapters by distinguished contributors.

"Full of information invaluable to the student, the house surgeon, and the general practitioner."—*Lancet*.

Third Edition. Demy 8vo. 79 Illustrations, 44 in Colours. 12s. 6d. net. Postage 9d.

## **DISEASES OF THE THROAT, NOSE & EAR**

For Practitioners and Students.

By the late **W. G. PORTER, M.B., B.Sc., F.R.C.S. Ed.**

Fully revised under the Editorship of **A. LOGAN TURNER, M.D. Ed., M.B., C.M., F.R.C.S. Ed.**

Consulting Surgeon, Edinburgh Eye, Ear and Throat Infirmary, etc.

AND OTHER COLLABORATORS.

"We know of no manual of its size better suited to the needs of the general practitioner."—*British Medical Journal*.

"A source of sound teaching of moderate dimensions for the senior student and general practitioner."—*Lancet*.

**Bristol: JOHN WRIGHT & SONS, LTD.**

London: **SIMPKIN & CO., Ltd.**

The Tonic Food richest  
in **VITAMINS**, and  
therefore of great value in  
all wasting diseases. Delicate  
children thrive on it.

The  
"A.B.C."  
of Diet.

# **'ROBOLEINE'**

Samples and Literature on application to  
**OPPENHEIMER SON & CO., LTD.,**  
179, Queen Victoria Street, London, E.C. 4

## **PENDYFFRYN HALL SANATORIUM**

(Nordrech-in-Wales.)

For the Open-air and Inoculation Treatment of all Forms  
of Tuberculosis.

**TREATMENT AVAILABLE BY ARTIFICIAL PNEUMOTHORAX.**

One of the first Sanatoria opened in the United Kingdom to carry out the treatment as practised at Nordrach. Carefully graduated walks rise through pine, gorse, and heather to a height of over 1,000 feet above sea level, commanding extensive views of both sea and mountains. Sheltered from E. and N.E. winds. Climate mild and bracing. Small rainfall. Large average of sunshine. There are over five miles of walks in the private grounds. Rooms heated by hot-water radiators and lit by electric light. Sister and full Nursing Staff. Trained nurse on duty all night.

Resident Medical Officers: **R. ARDRA FEGAN, M.R.C.S., L.R.C.P., L.M.S.S.A.; F. A. LUCAS HAMMOND, M.D., M.R.C.P., D.P.H.**

For particulars apply to the Secretary, Pendyffryn Hall, Penmaenmawr, North Wales.

Telegrams: Pendyffryn. Telephone: 20 Penmaenmawr.

**ILLUSTRATED BOOKLET FORWARDED ON REQUEST.**

## BOOKS.

---

### THIS MONTH'S SPECIAL REVIEWS.

---

*Le Bactériophage son rôle dans l'immunité.* By F. d'HERELLE.  
Paris, Masson et Cie. 12 Fr.

VIOLENT shocks often do harm but occasionally do good, and this work is one we would recommend to any complacent and self-satisfied bacteriologist—if indeed such a person exists—who believes that bacteriology and immunity have reached finality. The shock of reading this book should administer a much needed corrective to his equanimity. We would further recommend the book to all those who have any interest in the science of medicine and who realise that text-books are not the last word but are merely necessary, though unsatisfactory, substitutes for monographs by authors who are experts in the subjects dealt with by them.

In making these recommendations we are not to be taken as agreeing with everything that the author states: in some respects we most emphatically disagree, but the careful reading of d'Herelle's experiments and conclusions cannot fail to make one pause and consider certain subjects which, up to the present, have either been unknown or have aroused no particular interest.

"Le Bactériophage" is a monograph based on an enormous mass of experimental work done by the author and his fellow-workers. The greater portion of this work has already been published in various journals, but here one can, for the first time, get a connected account of d'Herelle's views on the bacteriophage.

Competent bacteriologists disagree with many of the author's conclusions, but as regards the fundamental phenomena, which are of the most novel and surprising nature, every observer who has devoted a little time and trouble to the work has been forced—even against his pre-formed opinions—to agree that d'Herelle's findings are easily capable of reproduction. The reviewer has obtained results on which d'Herelle's written account and illustrations might have been based.

d'Herelle's interest was first aroused by the following experiment. He took a small amount of a stool of a patient convalescent from dysentery (Shiga type) and emulsified it in saline. Next day the saline suspension was filtered through a porcelain filter which retained all ordinary bacteria. A small amount of the filtrate was added to a broth culture of Shiga's bacillus. In a few hours this broth culture had become clear, and, when attempts were made to obtain sub-cultures, it was found that all the bacteria were dead. If this lysed culture was, in turn, filtered through a porcelain candle and a few drops of the filtrate added to a new broth culture of the same bacillus, clearing of the latter occurred. This phenomenon could be reproduced, apparently indefinitely, in series. So far from becoming weaker by these successive passages the substance which produced the solution of the bacteria became stronger, solution occurring more quickly than with the first filtrate obtained from the patient's fæces. This primary observation was repeated many times by d'Herelle and later by bacteriologists in almost every country. Whatever the explanation may be, there is no doubt about the facts.

The next step was also of great interest. Active filtrate was added to a broth culture and the mixture incubated. At intervals of one hour one drop was removed and spread on the surface of agar. These sub-cultures were incubated and examined the next day. The first showed a uniform culture of the bacillus except for two circular spots, about 2 m.m. in diameter, which were apparently bare. The second tube showed 6 similar spots, the third 100, while the fourth had no visible growth of bacteria. By this technique it has been possible to demonstrate in fæcal filtrates a solution-causing substance when this was too feeble to cause complete lysis and clearing of all the bacteria in a broth culture. If, on spreading on the surface of agar, clear spaces were seen, the substance was present and, by successive passages through a series of broth cultures, it was possible to exalt its activity so that ultimately it was able to clear completely suspensions of bacteria. With this technique one is now able to examine stools, and to detect in them the presence of the solution-causing substance with ease and certainty. Such substances

# Essence of Rennet Concentrated

(Benger). The highest quality sweet essence, for professional use in Infant and Invalid Feeding.



Whey for diluting the milk for young infants prepared with Benger's Essence of Rennet is of the greatest value in the treatment of diarrhoea, vomiting, etc., and can be used with perfect safety and good results. Makes splendid junket.

*Benger's Essence of Rennet was specially prepared for H. Ashby, M.D., for producing whey in Infant Feeding (see his books).*

BENGER'S FOOD, Ltd., — MANCHESTER, England.

Branch Offices—NEW YORK: 99, Beekman Street.

SYDNEY: 117, Pitt Street.



I.116

## *Hemaboloids* ARSENIATED (with Strychnia)

furnishes the Tonic Trinity, comprising Iron, Arsenic and Strychnia, in permanent and palatable solution, reinforced by Predigested Albuminoids (Peptones) and Bone Marrow Extract.

Each Tablespoonful represents the equivalent of

20 Minims r. Ferri Chlor.  
1/40 Grain Arsenious Acid.  
1/80 Grain Strychnia.

*Furnished in two sizes at 2/3 and 4/6.*

JOHN MORGAN RICHARDS & SONS, LTD.  
46 HOLBORN VIADUCT—LONDON E.C.



*Introducing*

## Iron in Large Doses

**T**HE theory of the treatment of Anæmia, Chlorosis, General Debility, etc., by the application of Iron in Large Doses is now well known, experience having proved conclusively that the small doses of Iron usually prescribed hitherto are slow and unreliable, and that anæmic conditions react much more successfully to Large Doses.



## IDOZAN

IDOZAN is a neutral, permanent, easily assimilable form of Iron, containing 5 per cent. Iron as Ferric Hydroxide in a labile colloidal solution, a small quantity of sugar being added as protective to the colloid.

Its sweetish aromatic taste makes it pleasing to the palate, and it is entirely free from that astringency usually found in Iron preparations, especially when the Iron is present in large proportions. Its liquid form ensures efficient absorption, and it does not irritate the stomach or intestinal canal. As it has no constipating effect, there is no need to add the usual laxative for correction.

*One teaspoonful of Idozan contains about 4 grains pure iron ; a dessertspoonful about 8 grains ; and a tablespoonful about 16 grains.*

*Idozan is sold only in bottles of 8 ozs., and may be ordered from any Chemist, but for Hospital use bottles containing about 2 lbs. are supplied. ————— SAMPLES ON REQUEST.*

**Chas. Zimmermann & Co. (Chem.), Ltd., Ferrosan Dept.  
9-10 St. Mary-at-Hill, London, E.C.3.**



have been found in the stools both of the healthy and of those suffering from many diseases. The substance may be active against one or more types of bacteria. The anti-Shiga substance is the one most easily demonstrated but those active against the other dysentery bacilli, against members of the typhoid group or against other organisms have been found. The activity is not necessarily limited to one type of organism. In one found by the reviewer, for example, the substance isolated was very active against *B. dysenteriae* (Shiga), moderately against Flexner and *Y. dysentery* bacilli, very slightly against *B. paratyphosus* B, and not at all against *B. coli*, *B. typhosus* and *B. paratyphosus* A.

The explanation of this lysis of bacteria is exceedingly difficult. Lysis might be effected by—1, some substance found in the intestine; 2, autolysins formed by the bacteria under the influence first of all of some intestinal substance; or 3, by a living organism which thrives at the expense of the bacteria. The first view is untenable for the lysin has been found active after 1,500 passages, the amount transferred each time being  $\frac{1}{1000}$  c.c. Many observers incline to the second opinion, but d'Herelle believes implicitly in the third, and has given the name of *Bacteriophagum intestinale* to the organism. In either case the difficulties are great. Any conservative bacteriologist must, at first, be disinclined to believe in the possibility of a living organism so small as to be able to act as a parasite on bacteria and yet so highly specialised in function, but we must remember that occasionally the truth is highly improbable. d'Herelle's opponents also have great difficulty in making their theory fit in with all the observed facts, and yet it is hard to imagine any other possible solution. d'Herelle's chief grounds for his belief are the following:—1, the appearance on agar of clear spaces indicating that the substance acts as a solid body, not diffused throughout the medium in solution. 2, his observation of minute granules which he describes as penetrating and multiplying in the bacterial body. 3, the fact that the activity of the filtrate is destroyed by heat and by various disinfectants not inhibiting other enzymes. 4, no increase in activity and no solution of the bacteria occurs unless the bacteria are living. 5, the activity of a filtrate acting on a culture increases in

waves, the period being about one to one and a half hours. This he explains as the period necessary for one bacteriophage to increase in a bacterium to 15 to 25 individuals when the bacterium disrupts, liberating the young bacteriophages. 6, the possibility of "educating" the substance to act on other species of bacteria—an exclusive property, according to d'Herelle, of living matter. 7, various rather complicated experiments in complement fixation and immunity reactions.

The reviewer refuses to commit himself for or against d'Herelle's theory of the existence of this obligatory bacterial parasite, the *Bacteriophagum intestinale*. d'Herelle's writing is convincing and his logic apparently sound, but the time is not yet ripe for a definite opinion.

For d'Herelle the bacteriophage is the most important factor in immunity. Just as bacteria wage war against man, so the bacteriophage wages war against bacteria. The bacteriophage may have its virulence exalted or attenuated, the bacteria may be conquered or, by producing a more resistant type, be victorious. "*L'histoire de la maladie est en définitive l'histoire de la lutte entre le Bactériophage et une bactérie.*" "*En un mot, la guérison se produit toujours par suite d'une exaltation de la virulence du Bactériophage, suffisante pour lui permettre de parasiter et de détruire les bactéries pathogènes implantées dans l'organisme. La mort survient, soit par suite de l'inertie du Bactériophage soit par suite de l'acquisition par la bactérie de l'état réfractaire, ce qui permet à cette dernière, dans l'un et l'autre cas, de se développer sans entraves.*"

d'Herelle is, as one would expect, optimistic as to the utility of his bacteriophage both therapeutically and prophylactically, but one must confess that the accounts which he gives of cases treated by "bacteriophage-therapy" are less numerous and less convincing than are the accounts of his "in vitro" experiments. A given bacteriophage can, however, be administered to a man by mouth and be isolated subsequently from the stools, so possibly it may have a future in medicine. Let us hope that it will not be brought into the disrepute which has followed the commercialisation of vaccine therapy.

d'Herelle's outlook is wide, and he predicts the day of wholesale immunity conferred by the bacteriophage. "*Nous*

pouvons donc envisager la possibilité de l'immunisation collective de la population car il serait facile de mélanger à l'eau potable des cultures du Bactériophage, surtout dans les agglomérations urbaines où l'eau est distribuée par canalisation."

Finally, let us recommend once more this book as one to be read carefully and thought over. Even to those disagreeing completely with d'Herelle's conclusions it will be found of great interest.

J.W.B.

*Therapeutic Immunisation.* By W. FORD ROBERTSON.  
Edinburgh : E. and S. Livingstone.

THE author of this book reminds us of the character in "Alice in Wonderland" who would not be ruled by mere words, but who ruled them instead and made them mean exactly what he desired. For this reason the work tends to confuse the reader who must, before understanding it, learn the terminology. As an example we may give the word "hæmolysis" as used in the classification of streptococci. For Robertson, hæmolysis does not mean the liberation of hæmoglobin from red blood cells, but something quite different. So streptococcus pyogenes is either hæmolytic or non-hæmolytic but the pneumococcus is always hæmolytic. Robertson recognises not only the four classical strains of pneumococci but also the "pneumococcus of pernicious anæmia" and the "pneumococcus of rheumatoid arthritis."

Those who have some experience of vaccine therapy or rather of "therapeutic immunisation" may become jealous on reading of the author's many successes. Some may even be sceptical. Gastric and duodenal ulcers, appendicitis, chronic intestinal stasis, neuralgia, neuritis, neurasthenia, exophthalmic goitre, disseminated sclerosis, and various acute insanities, to mention but a few, are cured, or at least improved, by "therapeutic immunisation." Staphylococcus aureus vaccine is a specific for chilblains and B. coli vaccine for varicose ulcers. These last two are, however, examples of the pharmaceutical and not the specific action of vaccines. If the local disease yields no organisms, satisfactory growths of suitable bacteria may be obtained from the nose, mouth,

or throat, from the urine, or from the stools. "Anærobic diphtheroid bacilli" and "Streptococcus fæcalis hæmolyticus" are particularly potent bacteria in the author's hands.

We cannot recommend this work for general reading by the profession. One reason is that the clinicians might become dissatisfied with their present bacteriologists. There are also other reasons.

*Preventive Medicine and Hygiene.* By MILTON J. ROSENAU.  
3rd Edition. New York and London. D. Appleton and Co. 45s. net.

We have nothing but praise for this work, and congratulations for its author. A book of 1,567 pages which costs 45s. is not one which will be purchased by many without good reason. Yet three editions of this book have been exhausted, and the present is the fourth. The medical public have evidently reviewed this book, and have thoroughly approved of it. We do not profess to have read through this new edition, but we have carefully examined several of its sections and have found them clear, thoroughly revised and modern. It is not a student's book, but to any doctor who is concerned in any aspect of preventive medicine or hygiene it will be invaluable. The paper, printing, diagrams and illustrations are above criticism.

*Colloid Chemistry of the Proteins.* WOLFGANG PAULI. J. and A. Churchill. 1922.

THIS small book represents Part I. of a work by the director of the Physico-Chemical Biology Laboratory in the University of Vienna. It has been primarily derived from a course of lectures delivered in 1912-13, and represents much of the work done on the colloid proteins by Pauli and his associates. Designed to meet the requirements of the chemical physiologist it naturally offers little inducement to the dilettante student who wishes to keep himself level with the trend of modern research and achievement in this particular branch. The iso-electric reaction is fully dealt with in relation to the proteins. Full tables of results with accompanied detailed calculations and curves are given.

*Intrinsic Cancer of the Larynx and the Operation of Laryngo-Fissure.*" By IRWIN MOORE. University of London Press, 1921.

THE operation of laryngo-fissure has contrived to itself an interesting and unusual history. First suggested so long ago as 1812, it was not actually performed until 1833, and during its earliest years gave promise of great success. Later on, however, owing to faulty technique and the choice of unsuitable cases, it proved disappointing and suffered disrepute until Durham and Butlin re-established its value by greater care in execution and pre-operative diagnosis. Since then, a steady increase in successes has eventually established laryngo-fissure as one of the bright spots in the treatment of cancer. Such being the position to-day, one eagerly looks forward to the publication of a book which deals with so important an operation from all points of view. In Dr. Irwin Moore's modest sized volume we have all that there is to be known up to the present about laryngo-fissure. A most cordial welcome should be extended to a book which carefully deals with all the practical points before and after the performance of the operation. Not the least interesting is the historical section, which deals fully and concisely with all the early attempts to open the larynx for intrinsic growths.

One point might be specially referred to—the question of anæsthetic. In the author's opinion preference would appear to be given to chloroform. If we might suggest an omission in such an excellent book, it is that the merits of ether given by the colonic method have not been placed on record. In several cases of laryngo-fissure recently performed both for foreign bodies and growths, we have found this a method of preference allowing of an easy and peaceful narcosis, while reducing the duties of the anæsthetist near the operation field, to the occasional administration of a few drops of ether by way of the tracheotomy tube—undoubtedly a distinct benefit where fine work needs all the room possible.

We would again extend our cordial welcome to this monograph, which should be in the hands of all laryngologists, and which must be looked upon as one of the most valuable contributions to the surgery of the larynx which has appeared for many years.

J.S.J.



*Essays on Surgical Subjects.* By SIR BERKELY MOYNIHAN.  
W. B. Saunders and Co., 1921.

THIS is one of those books which unopened awakens keen anticipation. One counts beforehand upon a style that will be lucid and epigrammatic, a style in fact which has helped to shift the centre of English surgery to Leeds. These essays include a tribute to Murphy, "the greatest clinical teacher of his day," which is written with an enthusiasm that gives the reader a sense of personal loss. This is followed by the "Ritual of a Surgical Operation," a title alone worth much to those who teach surgery. If the master discards gloves, the pupil may neglect soap. Ritual ingrains reverence.

Dealing with gastric ulcer Moynihan states that no diagnosis of chronic ulcer should now be confidently accepted unless the ulcer is revealed by x-ray examination, or is displayed upon the operation table.

Other essays treat of "Intestinal Stasis,"—a temperate account of a subject that has suffered from over-statement. "Acute Emergencies of Abdominal Disease"; "The Gifts of Surgery to Medicine" in which insistence is laid upon the pathology of the living; "The Surgery of the Chest" a valuable summary of personal experience and European teaching.

These essays, like the plays of Wilde, should be read separately. Iteration dulls the flash of epigram.

*Clinical and Operative Gynæcology.* By PROFESSOR J. M. MUNRO KERR, M.D., F.R.C.P.S. Professor of Obstetrics and Gynæcology, Glasgow University, etc. Cr. Svo. Pp. 832 + XVI. London: Henry Froude and Hodder and Stoughton, 1922.

It is a well-known fact that many great actors have only one part in which they are really great. Professor Munro Kerr has written a very great work which will last: it is not the volume at present under review—it is his operative midwifery. We took up *Clinical and Operative Gynæcology* with a gladsome heart—we put it down with a sigh—not of content, but with a certain degree of grief, for the book just fails to get home.



Although we know and appreciate the author's worth, his great experience would have been more in evidence had the style been more impersonal. The work is divided into three parts (1) Clinical Gynæcology, 560 pages. (2) Medical Treatment and Remedies, 18 pages. (3) Operative Gynæcology, about 200 pages. There are valuable special articles on development, nervous disorders, venereal diseases, anæsthesia and transfusion. We are not sure that the arrangement of certain points simplifies the comprehensiveness of the text, *e.g.*, there is no description of the uterine ligaments in the anatomy section, but we are referred to the section on displacements; and in dealing with tuberculosis of the tubes we are referred back to the chapter on general tuberculosis. Voluminous references in a work of this kind are unnecessary, but we are surprised to find no reference to the work of Frankl and others on the corpus luteum. A classification of ovarian cysts would have been valuable. The paragraphs on sterility will be improved in succeeding editions, for at present they seem to be incomplete. The author suggests a very simple description of the supports of the uterus and vagina. He divides these supports into four tiers (1) Levator-coccygeus muscles. (2) Paracervical tissue. (3) Uterine ligaments. (4) Cervical muscles. Nothing could be better than this.

At some utopian day authors of text books in gynæcology will give a clear definition of the normal position of the uterus. The student is expected to know this, and yet it is either omitted as here, or given vaguely as by many authors. We are amazed to find that the dangerous old-fashioned septic method of replacing the uterus with a sound is recommended, described, and illustrated by two figures.

Another point which requires more agreement is the definition of a tumour of the vulva; the author puts under this heading femoral hernia, hæmatoma vulvæ, varicose veins; many professors will refuse to accept these conditions as tumours.

There is a thoroughly sound and comprehensive article on chronic endometritis which explains logically the old and the modern ideas. We fear that the treatment of acute salpingitis will not be clear to the puzzled practitioner, but the

chronic condition is explained in a manner which could not be excelled.

"We are perfectly justified in discussing this complication under a special heading." Yes, perimetritis and parametritis might be discussed under a separate heading if a convincing differential diagnosis could be given, but it is not given here, and we are sceptical if it can be given at all. It may be thought that we have criticised some points in this work too severely, but a gynæcologist of the standing of Professor Munro Kerr must stand, and is well able to stand criticism. In the last part operations are dealt with magnificently; they are described more in the form of an atlas than anything else. Most known operations are considered, and this section alone makes *Clinical and Operative Gynæcology* a valuable addition to the library of the gynæcologist.

B.S.

*The Transactions of the Edinburgh Obstetrical Society.* Vol. XVI. Session 1920-21. Edinburgh: Oliver and Boyd 1922. Pp. 148 + XXVII.

As a record of work done, this volume compares favourably with past records, but we regret that our plea when reviewing the 39th volume that cross-compliments be omitted has not been acceded to. We consider that the blue pencil of the editor should strike out the numerous congratulations which are scattered through the various valuable discussions.

Probably the contribution which will appeal most to the reader is that of Dr. Haultain on "Twilight Sleep"; this well-known writer has now had a very large experience of the method, and gives it his unqualified approval—a sentiment which was shared by the speakers on the paper. Eclampsia which might be called the fashionable obstetric subject at present has two papers, one in its relation to concealed accidental hæmorrhage, the other to placenta prævia.

Dr. Young's communication on an organism obtained from carcinomatous growths is republished in these transactions, and there is a useful contribution to the literature of uterine prolapse.

B.S.

*A Short Practice of Midwifery for Nurses.* By HENRY JELLETT, M.D., F.R.C.P.I. Gynæcological Surgeon Canterbury Hospital, New Zealand, etc. Pp. 427 + XIII. Sixth edition revised. London: J. H. Churchill, 1922. Pp. 427, XII.

DR. JELLETT has emigrated to New Zealand since the last edition of this popular work was published. The present revised edition is up to the standard of its predecessors, and some necessary alterations and additions have been made. We are confident that the well-deserved popularity of the previous issues will be maintained.

B.S.

---

# ABSTRACTS OF CURRENT LITERATURE.

## GYNÆCOLOGY AND OBSTETRICS.

LEE, GEO. : *The Test of Labour : Results in 100 Consecutive Deliveries.*

"Surg. Gyn. Obst. XXXV. I. 63."

DANFORTH : "*Is Conservative Obstetrics to be Abandoned ?*" Amer. Jour. Obst. and Gyn., III. 6. 609.

WE believe and hope that the boom in surgical obstetrics has reached its zenith. It is satisfactory to find two of the premier journals in the obstetrical world containing pages on this subject. Lee makes a very strong plea for a test of labour. He believes that this is obtained by allowing labour to proceed, until childbirth either result spontaneously, or from operative assistance that reinforces the natural powers so that birth is through the natural passages. Of course, when there are obvious obstacles to delivery no patient should be submitted to the test. The paper shows most satisfactory results in an analysis of 100 consecutive cases.

Danforth asks "Is conservative obstetrics to be abandoned ?" He analyses 500 cases and proves the efficacy of conservatism to the hilt. Among other points he deals with the Potter elective version and demonstrates that his (Danforth's) foetal mortality is 1.4 per cent. as compared with 7.5 per cent with the Potter method. These two papers are published about the same time. Obviously America is recovering from the operative disease which was never very pronounced in Ireland.

BETHEL SOLOMONS.

HORNER, D. : *Röntgenography in Obstetrics.* "Surg., Gyn. Obs. XXXV. I. 67."

THIS paper suggests that *x*-rays may be used in obstetrics for (1) diagnosing early pregnancy. (2) foetometry before induction of labour, and to diagnose monsters before performing Cæsareansection. (3) pelvic deformities and measurements. (4) injuries to the newborn. He states there is no danger to mother or child. Davis (p.119 *ibid.*) remarked that he frequently found it necessary to have teeth *x*-rayed in pregnancy, thus avoiding the results of toxæmia from septic dental foci.

BETHEL SOLOMONS.

# NEURASTHENIA

The tumultuous life entails its penalties—among them neurasthenia. The storm and stress of modern civilization exhaust the reserve force of the organism just as the suffering and hardships of war did. Organotherapy is effective in overcoming the nervous exhaustion induced more subtly, but just as surely, by the high speed conditions of the Twentieth Century.

## HORMOTONE

tends to restore the perfect hormone balance essential to the maintenance of health and to the restoration of it in a run down condition.

In neurasthenic cases associated with a high blood pressure use

### **Hormotone Without Post-Pituitary**

Dose of either preparation: One or two tablets three times daily before meals.

**G. W. CARNRICK CO.**

**417-421 Canal St.,**

**New York, U.S.A.**

*Distributors :—*

**THE AMERICAN DRUG SUPPLY CO., 1 CHARING CROSS,  
LONDON, S.W.1.**

The combination of tonics and stimulants explains the clinical results obtained in the treatment of nervous disorders by the use of

## FELLOWS' COMPOUND SYRUP OF HYPOPHOSPHITES

*"A true stabilizer of shaken nerves"*

It contains the "mineral foods", Sodium, Potassium, Calcium, Manganese, Iron and Phosphorous, and the stimulating agents, Quinine and Strychnine.

*Samples and Literature sent upon request.*

**FELLOWS MEDICAL MANUFACTURING CO., Inc.**

26 Christopher Street, New York, N. Y.

FOR INTERNAL TREATMENT OF GONORRHOEA,  
URETHRITIS AND OTHER AFFECTIONS OF THE  
GENITO-URINARY TRACT.

### Santal Midy Capsules

have been prescribed with uniform success for over 30 years. Distilled from carefully selected Mysore Sandal Wood, the oil is bland and remarkably  
**FREE FROM THE IRRITANT AND NAUSEATING EFFECTS**  
which are provoked by many preparations.

# SANTAL MIDY

There is marked absence of Gastric and other disturbances, diarrhoea and skin eruptions. Its mild chemotactic properties permits its administration in relatively large doses without fear of too violent reaction or intolerance.

### SANTAL MIDY CAPSULES

may be prescribed and relied upon in all stages of Gonorrhoea and in other forms of Urethritis and affections of the Genito-Urinary tract. The Capsules contain 5 drops, and usually 10 to 12 are given daily in divided doses.

*Prepared in the Laboratoire de Pharmacologie Generale,  
8 Rue Vivienne, Paris, and sold by most Chemists and Wholesale Druggists throughout the world.*

U. K. Agents—Wilco, Gozeau & Co., 15 Gt. St. Andrew St., W.C. 2



KENNEDY, RONGY AND ROSENFELD: *Sterility*. "Amer. Jour. Obst. and Gyn. 111. 6. 607." "Amer. Jour. Obst. and Gyn. 111. 6. 496."

WE are glad to find that our work here has induced gynæcologists in America to try and find a suitable method for keeping the tubes open after resection. Kennedy uses Cargile membrane instead of catgut which is our choice; in order to direct it through the tubes he suggests a hollow silver probe through which some piano wire is directed and draws the membrane after it; the method is best understood by the illustration which accompanies the article. In a paper by Rongy and Rosenfeld a plea is made for transuterine insufflation of the tubes with  $\text{CO}_2$ . The technique is given and makes interesting reading, but we believe there is less danger to the patient in performing laparotomy in doubtful cases. In a series of 100 patients there were no deaths, but the following complications occurred (1) syncope in an obese patient. (2) a second case of syncope. (3) an acute inflammatory condition in the left fornix.

BETHEL SOLOMONS.

PARDEE: *Cardiac Failure in Pregnancy*. "Amer. Jour. Obst. and Gyn. 111. 6. 496."

PARDEE considers that aortic regurgitation is more dangerous than mitral stenosis, but is more concerned with the symptoms or signs of failure than with the nature of the disease. Morphine with atropine is advised in labour; in addition, phlebotomy and digitoxin gr.  $\frac{1}{50}$  intravenously are recommended. Ether preceded by a small amount of chloroform is the best anæsthetic. If decompensation should occur in labour it is necessary to do Cæsarean section unless forceps can be applied. The treatment is the same in pregnancy except that oxygen may be given to relieve cyanosis.

BETHEL SOLOMONS.

RILEY: *An Unusual Case of Extrauterine Pregnancy*. "Amer. Jour. Obst. and Gyn. 111. 6. 630.

RILEY records notes of a most unusual case. The woman gave the history of a ruptured tubal pregnancy which was not operated on. After this she had four full-term children. Then she consulted Riley who examined, and decided that she had a myoma of the uterus. At operation the mass proved to be the bones of a foetus and much foul-smelling material. On assembly of the bones, the skeleton was a foetus about  $7\frac{1}{2}$  months. The patient recovered after a stormy convalescence.

BETHEL SOLOMONS.

DAVIS: *Neoplasia of the Kidney*. "Amer. Jour. Obst. and Gyn. 111. 5, 478."

THIS paper reports five primary renal tumours which have been carefully studied to exclude metastatic origin. They consist of (1) Papillary epithelioma. (2) Hypernephroma. (3) Malignant teratoma. (4) Squamous celled carcinoma. (5) Lympho-blastoma. Davis states that the developmental history of the renal tissues is yet incomplete, and at many points theoretical. The diagnostic symptomatology is often indefinite. Clinical and pathological investigation of renal tumours should be carefully made and reported.

BETHEL SOLOMONS.

## PHYSIOLOGY.

### *Recent Experiments on the Pituitary\**

SINCE Paulesco first, followed by Cushing, removed the entire pituitary in animals, it has been held that such removal results in death. These operators used the temporal approach. Later Aschner, using the buccal route, showed that pituitary extirpation is not necessarily fatal. Camus and Roussy have confirmed this finding, verifying the complete removal of the gland by histological examination. Where death ensued, it was due as a rule to hæmorrhage or meningitis.

It has been suggested that survival is explained by the presence of aberrant pituitary substance, yet this work seems to prove that as a differentiated organ the hypophysis is not essential to life.

### *Polyuria.*

Camus and Roussy have further shown that the polyuria which follows many pituitary operations in animals is not due to a lesion of the pituitary itself, but to injury to the part of the brain to which the gland is attached. This area, the tuber cinereum, is a mass of grey matter in the floor of the third ventricle, between the optic chiasma in front, and the corpora mammillaria behind.

At the centre of the tuber is a funnel-like projection, the infundibulum, which narrows to form the pituitary stalk. Spiegel, Zweig and Lhermitte have recently described four paired nuclei in this area; one above the chiasma, one more laterally placed, the supraoptic nucleus, and external to this, the nucleus proprius of the tuber. Lastly, a paraventricular nucleus which lies between the ependymal lining of the third ventricle and the anterior pillar of the fornix. These nuclei are connected with the pons and globus pallidus.

\*IIIe Réunion Neurologique Internationale Anuelle, Paris, June 3rd and 4th, 1922.

Camus and Roussy found that permanent diabetes insipidus occurred in a dog in which the pituitary was unharmed, but where experimental puncture had injured the nucleus proprius of the tuber. In another experimental animal with polyuria of five months' duration and an adiposo-genital syndrome, the pituitary was intact, but the two nuclei of the tuber and the paraventricular nuclei had been injured.

A further series of experiments showed that when the entire pituitary was removed, no polyuria resulted unless the region of the tuber was involved, nor did any puncture remote from the tuber produce polyuria. The paired nucleus proprius of the tuber is very superficial, and thus polyuria may follow a superficial injury; an animal of five kilograms weight may, for example, pass three litres of urine daily during many months.

Glycosuria was only rarely produced in these experiments; when it occurred it was thought to be due to a basal rather than to a pituitary lesion. It is noteworthy too, that hydro-carbon tolerance was not affected by lesions limited to the gland or by its complete removal. Hanchett, *American Jour. Med. Science*, May, 1922, p. 685, describes an independent series of experiments along similar lines. These include stimulation of both lobes of the gland directly with a heated rod, and indirectly, by filling the sphenoidal sinuses with hot wax. In only one animal of this series did polyuria occur, and in this the cautery had failed to reach the pituitary, but had injured the corpora mammillaria. Electrical stimulation with induced currents produced polyuria, but as it is difficult to limit their diffusion, the urinary increase cannot with certainty be set down to stimulation of the gland.

In every case where there was traction upon the pituitary stalk, polyuria ensued. Cushing has observed that the incidence of polyuria was greater in operated pups than in adult dogs, and Hanchett suggests that, with a temporal approach, the upward displacement of the brain required to secure effective exposure in pups must be relatively greater, and involve more traction upon the pituitary stalk. Stalk-traction thus seems to explain the "unaccountable postoperative polyuria of extreme grade" in a number of Cushing's cases. Hanchett's work thus confirms in a general way that of Camus and Roussy, which is also supported by the experiments of Bailey and Bremer in Belgium. Hanchett, however, finds that cauterisation of the mammillary bodies produces polyuria, while Camus and Roussy restrict the polyuric area to the region of the tuber.

These findings already receive a measure of clinical confirmation, though few pathological reports have been sufficiently discriminating to bear useful testimony regarding these precise experiments. Hagenbach in 1882 described a case of tuberculosis meningitis with polyuria, in which a caseous nodule was found in the infundibulum, associated with dilatation of the fourth and lateral ventricles. Lhermitte has described a case of diabetes insipidus in which there

was a syphilitic lesion of the nuclei of the tuber, the pituitary being unaffected. Others have described lesions, with polyuria, varying from complete destruction of the pituitary to disease of the stalk alone. Simmonds collected ten cases metastatic tumours of the pituitary, only three of which showed polyuria. In one case of this series *without* polyuria the entire gland was destroyed, except for a small piece of the anterior lobe; in another, however, with normal excretion the pars nervosa, the pars intermedia and the infundibulum had been replaced by the tumour. Demole of Geneva, on the other hand, reports polyuria in a case of dementia precox with an adiposo-genital syndrome, which showed no lesion of the pituitary nor of the tuber, but in which the interstitial cells of the testis were definitely altered.

Earlier workers have stated that polyuria can be produced by the stimulation of other regions of the brain than that of the pituitary. Claude Bernard in 1849 secured an increased output of urine by injuring the floor of the fourth ventricle in front of the spot for glycosuric puncture. Kahler produced in rabbits a lasting polyuria, with polydipsia, by injecting concentrated silver nitrate solution in small quantity into the corpus trapezoides of the pons, and into the lateral part of the medulla. (The connections of pons with the nuclei of the tuber are suggestive in regard to this.)

#### *The Adiposo-genital Syndrome.*

This syndrome has hitherto been considered as a manifestation of hypopituitarism but Camus and Roussy find that when the pituitary is removed from dogs without injury to the base of the brain no genital atrophy, obesity, nor increased hydrocarbon tolerance occur. A pure lesion of the base, however, may produce an increase in weight of seven kilograms in three weeks.

---

# LIST OF PERIODICALS AND PUBLICATIONS RECEIVED IN EXCHANGE FOR "IRISH JOURNAL OF MEDICAL SCIENCE,"

## AMERICA :—

*Southern Medicine and Surgery.*  
*The Medical Record.*  
*The Medical Card.*  
*The American Journal of Insanity.*  
*The New York Medical Journal.*  
*The Journal of the American Medical Association.*  
*The Archives of Pediatrics.*  
*The John Hopkins Hospital Reports.*  
*Cohio State University College of Medicine.*  
*Asclepios Habana.*  
*College of Physicians of Philadelphia.*  
*New York Academy of Medicine.*  
*American Journal of Medical Sciences.*  
*American Journal of Obstetrics and Gynæcology.*  
*American Journal of Syphilis.*  
*Archives of Diagnosis.*  
*Medical Clinics of North America.*

## AUSTRALASIA :—

*The Australasian Medical Gazette.*  
*Medical Journal of Australia.*  
*New Zealand Medical Journal.*

## BELGIUM :—

*Bruxelles Medical.*  
*Bulletin de Académie Royale Belgique.*

## CANADA :—

*Canadian Medical Journal.*  
*The Journal of the Canadian Medical Association.*

## DENMARK :—

*Hospitals-Tidende.*  
*Ugeskrift for Læger.*

## FRANCE :—

*Archives Générales de Chirurgie.*  
*Annales Médico-Psychologiques.*  
*Bulletin de l'Académie de Médecine.*  
*Gazette des Hôpitaux.*  
*La Presse Médicale.*

*Revue Hebdomadaire de Laryngologie.*

*Lyon Médical.*

*Journal de Médecine de Bordeaux.*

*Revue de Chirurgie.*

*Revue de Laryngologie, d'Otologie et de Rhinologie.*

GERMANY :—

*Zentralblatt für Chirurgie.*

GREAT BRITAIN :—

*Proceedings of the Royal Society of Medicine.*

*The Edinburgh Medical Journal.*

*The Lancet.*

*The British Medical Journal.*

*The Journal of Mental Science.*

*The Glasgow Medical Journal.*

*The Practitioner.*

*The Journal of Anatomy and Physiology.*

*The British Journal of Dermatology.*

*The British Journal of Children's Diseases.*

*The Liverpool Medico-Chirurgical Journal.*

*Guy's Hospital Report.*

*British Journal of Ophthalmology.*

*Quarterly Journal of Medicine.*

*The Hospital.*

*British Journal of Tuberculosis.*

*Medical Science Abstracts and Reviews.*

*The Journal of Neurology and Psychopathology.*

INDIA :—

*The Indian Medical Record.*

*Indian Medical Gazette.*

ITALY :—

*Lo Sperimentale.*

*Archivio di Pathologia e Clinica Medica.*

*Archiviodi orthopedia.*

SCANDINAVIA :—

*Acta Medica Scandinavica.*

*Acta Chirurgica Scandinavica.*

NORWAY :—

*Norsk Magazin for Laegevidenskaben.*

*Hygeia.*

SWEDEN :—


*Upsala Lakare forenings Forhandlingar.*

SWITZERLAND :—

*League of Red Cross Societies.*

*Revue Médicale de la Suisse Romande.*





"I have much pleasure in testifying to the superiority of Liquor Bismuthi 'Schacht' over all other preparations of the same kind; not alone from the fact that it does not deposit as other solutions of Bismuth do, but that it is the only one whose therapeutical action is entirely sure and satisfactory."

-M.D. Lond.

**BISMUTHUM CRYST.**

**ROYAL COLLEGE OF PHYSICIANS OF EDINBURGH.  
ROYAL COLLEGE OF SURGEONS OF EDINBURGH.  
ROYAL FACULTY OF PHYSICIANS AND SURGEONS OF GLASGOW.**

Copies of **REGULATIONS** for the **TRIPLE QUALIFICATION** (**L.R.C.P.E.**, **L.R.C.S.E.**, and **L.R.F.P. and S.Q.**) and the **DIPLOMA IN PUBLIC HEALTH**, containing dates of Professional Examinations for year, 1922-23, Curriculum, etc., may be had on application to:—

**Mr. D. L. EADIE**, 49 Lauriston Place, Edinburgh, or to  
**Mr. WALTER HURST**, 242 St. Vincent Street, Glasgow (Inspectors and Treasurers at Edinburgh and Glasgow respectively.)

**ROYAL COLLEGE OF SURGEONS OF EDINBURGH.**

(Incorporated 1505).

**COPIES OF REGULATIONS** for the **FELLOWSHIP LICENCE**, **HIGHER DENTAL DIPLOMA**, and **LICENCE** in **DENTAL SURGERY**, containing dates of Examinations, may be had on application to:—

**D. L. EADIE**,  
Clerk of the College,  
49 Lauriston Pl., Edinburgh.

**ROYAL COLLEGE OF PHYSICIANS  
OF EDINBURGH.**

THE EXAMINATIONS for the LICENCE of the College (as a single qualification) for the ensuing year will be held on the first Wednesday and the following days of every month (except September and October).

Candidates for the LICENCE (as a single qualification) must send their names to the Secretary One Week before the date of the Examination at which they propose to appear.

THE EXAMINATIONS for the MEMBERSHIP of the College are held on the second Tuesday and the following days of January, April, July, and October.

Candidates for the MEMBERSHIP must submit their Applications and Testimonials to the Secretary One Month before the date at which they wish to appear for Examination.

For the Regulations in regard to the various qualifications granted by the College, and all other information, application may be made to the Secretary.

**J. S. FLOWER, M.D.,**  
Hon. Secretary.

# **BAILLIÈRE, TINDALL & COX**

"UNIVERSITY SERIES" OF MANUALS.

## **JELLETT & MADILL'S MIDWIFERY**

THIRD EDITION

"The book is one of the fullest and best of the manuals on Midwifery in the English language."—*Edinburgh Medical Journal*.  
Pp. xiv + 1200. With 20 Plates and 570 other Illustrations in the Text. **42s. net**

## **MONRO'S MEDICINE**

FOURTH EDITION

"An exposition of medicine that is thoroughly in accord with the best teaching of the present day."—*The Lancet*.  
Pp. xxiv + 1045. With 47 Illustrations. Plain and Coloured. **21s. net.**

## **CASTELLANI & CHALMERS' TROPICAL MEDICINE**

THIRD EDITION

"The most comprehensive and up-to-date manual on the subject."—*Medical Press*.  
Pp. xii + 2436. With 16 Coloured Plates and 909 other Illustrations. **45s. net.**

## **ROSE & CARLESS' SURGERY**

TENTH EDITION

"There is no text-book of surgery in this or any other country so adequate and yet so convenient."—*Practitioner*.  
Pp. xii + 1560. With 18 Coloured plates and 600 other Illustrations. **30s. net.**

## **STEWART'S PHYSIOLOGY**

EIGHTH EDITION

"It is one of the very best text-books on the subject."—*The Lancet*.  
Pp. xxiv + 1132. With Coloured Plate and 467 other Illustrations. **21s. net.**

## **GREEN'S PATHOLOGY**

TWELFTH EDITION

"The student of all grades will find that this work will meet his wants."—*The Lancet*.  
Pp. x + 603. With 4 Coloured Plates and 243 other Illustrations. **22s. 6d. net.**

## **BUCHANAN'S ANATOMY**

FOURTH EDITION

**SYSTEMATIC and PRACTICAL, including EMBRYOLOGY, in ONE VOLUME. In the Old Nomenclature.** Edited by a COMMITTEE OF DEMONSTRATORS IN ANATOMY.

"As a text-book of anatomy for preparation it has no equal. For students we know of no better book."—*Middlesex Hospital*.  
Pp. xii + 1743. With 677 Illustrations, mostly Original and in Colours. **30s. net.**

**BAILLIÈRE, TINDALL & COX,**

8 HENRIETTA ST., COVENT GARDEN, LONDON, W.C.2.

Publishing Offices of "THE MEDICAL PRESS AND CIRCULAR."

# CASSELL'S MEDICAL BOOKS

## FOR STUDENTS.

### Clinical Methods

By **ROBT. HUTCHISON**, M.D., Edin., F.R.C.P.Lond., and **HARRY RAINY**, M.D., F.R.C.P. Ed., F.R.S.E. Seventh Edition. With 16 Colour Plates and 157 Illustrations in the Text. 12s. 6d. net.

### A Manual of Chemistry

Adapted to the Requirements of Students of Medicine.

By **ARTHUR P. LUFF**, M.D., B.Sc. Lond., F.R.C.P., F.I.C. and **HUGH C. H. CANDY**, B.A., B.Sc.Lond., F.I.C. Sixth Edition. Revised and Enlarged by **HUGH C. H. CANDY**. With 64 Illustrations. 12s. 6d. net.

### A Manual of Physics for Medical Students

By **HUGH C. H. CANDY**, B.A., B.Sc.Lond., F.I.C. (A Companion Vol. to Luff and Candy's Manual of Chemistry.) Second Edition, Enlarged. With a Colour Frontispiece and 280 Illustrations in the Text. 7s. 6d. net.

### Materia Medica and Therapeutics

By **J. MITCHELL BRUCE**, C.V.O., M.A., LL.D.Aberd., M.D., F.R.C.P. Lond., and **WALTER J. DILLING**, M.B., Ch.B.Aberd. Twelfth Edition, Revised. 10s. 6d. net.

### Herman's Difficult Labour

Revised and Enlarged by **CARLTON OLDFIELD**, M.D.Lond., F.R.C.S.Eng. Sixth Edition. With 198 Illustrations. 16s.

### Diseases of the Nose and Throat

By **Sir StCLAIR THOMSON**, M.D., F.R.C.P.Lond., F.R.C.S.Eng. Second Edition, Revised and Enlarged. With 12 Colour Plates, 10 Black-and-White Plates, and 337 Figures in the Text. 30s. net.

### Manson's Tropical Diseases

Seventh Edition, Edited by **PHILIP H. MASON-BAHR**, D.S.O., M.A., M.D. (D.T.M. and H.) Cantab., M.R.C.P.Lond. With 21 Colour and 6 Half-tone Plates, 404 Figures in the Text, and 31 Charts. 31s. 6d. net.

### Surgical Applied Anatomy

By **Sir FREDERICK TREVES**, Bart., G.C.V.O., C.B., LL.D., F.R.C.S., Eng. Seventh Edition. Revised by **Prof. Sir ARTHUR KEITH**, F.R.S., M.D., LL.D.Aberd., F.R.C.S. Eng., and **W. COLIN MACKENZIE**, M.D.Melb., F.R.C.S. Edin., F.R.S.E. With 153 Illustrations, including 74 in Colour. 12s. net.

### The Student's Handbook of Surgical Operations

By **Sir FREDERICK TREVES**, Bart., and **JONATHAN HUTCHISON**, F.R.C.S.Eng. Third Edition. With 162 Illustrations. 9s. net.

### The Student's Handbook of Gynæcology

By **G. ERNEST HERMAN**, M.B., F.R.C.P.Lond., F.R.C.S.Eng. Second Edition, Revised and Enlarged, with Additions by **R. DRUMMOND MAXWELL**, M.D.Lond F.R.C.S. Eng. With 6 Colour Plates and 194 Illustrations in the Text. 10s. net.

### Elements of Surgical Diagnosis

By **Sir ALFRED PEARCE GOULD**, K.C.V.O., M.S.Lond., F.R.C.S. Eng., and **ERIC PEARCE GOULD**, M.D., M.Ch.Oxon., F.R.C.S.Eng. Fifth Edition, Revised. With 16 Radiographic Plates. 12s. 6d. net.

### A Handbook of Midwifery

For Midwives, Maternity Nurses, and Obstetric Dressers.

By **COMYNS BERKELEY**, M.A., M.C., M.D.Cantab., F.R.C.P.Lond., M.R.C.S.Eng. Fifth Edition, Enlarged. With Colour Frontispiece and 74 Illustrations in the Text. 7s. 6d. net.

**CASSELL & CO., Ltd., La Belle Sauvage, London, E.C.4**



# Edinburgh Medical Series

TEXT-BOOKS FOR STUDENTS AND PRACTITIONERS

## A TEXT-BOOK OF GYNECOLOGY

By **JAMES YOUNG**, D.S.O., M.D., F.R.C.S. (Edin.).

Assistant Physician, Royal Maternity Hospital; Lecturer in Gynecology, Post-Graduate Course, Edinburgh University; University Clinical Tutor in Gynecology; Gynecologist, Provident Dispensary, Edinburgh, Examiner, Central Midwives Board for Scotland, etc.

Containing 181 Illustrations, some with colour introduced. Crown 8vo. Price 15s. net.

"We have no hesitation in saying that, for its size and scope, this is probably the best illustrated and most comprehensive text-book of gynecology at present available for the student or practitioner."—"Medical Officer."

## DISEASES OF THE EAR, NOSE, AND THROAT IN CHILDHOOD

By **DOUGLAS GUTHRIE**, M.D., F.R.C.S., Surgeon to the Ear and Throat Department, Royal Hospital for Sick Children, Edinburgh. Crown 8vo., 30 Illus. in the text. Price 5s. net.

## DISEASES AND INJURIES OF THE EYE

By **Wm. G. SYM**, M.D., F.R.C.S.E., Containing 25 full-page Illustrations (16 in colour). Crown 8vo. Price 12s. 6d. net.

## CEREBRO-SPINAL FEVER

The Etiology, Symptomatology, Diagnosis and Treatment of Epidemic Cerebro-Spinal Meningitis. By **CECIL WORSTER-DROUGHT**, M.A., M.B., and **ALEX. MILLS KENNEDY**, M.D. Demy 8vo., with 8 full-page Plates and 56 line Illustrations and Charts in the Text. Price 35s. net.

## DISEASES OF CHILDREN

By **A. DINGWALL-FORDYCE**, M.D., F.R.C.P.E. Second Edition. Containing 29 full-page Plates (8 in colour) and 84 Illustrations in the text. Crown 8vo. Price 18s. net.

## PRACTICAL PATHOLOGY, MORBID ANATOMY, AND POST-MORTEM TECHNIQUE

By **JAMES MILLER**, M.D., F.R.C.P.E. Containing 111 Illustrations and a Frontispiece in colour. Crown 8vo. Price 12s. 6d. net.

## MEDICAL APPLIED ANATOMY

By **T. B. JOHNSON**, M.B. Containing 146 Illustrations and 3 full-page Plates in colour. Crown 8vo. Price 12s. 6d. net.

## OTHER MEDICAL BOOKS.

By **W. G. AITCHISON ROBERTSON**, M.D., F.R.C.P.E.

**THE STUDENTS' GUIDE TO VACCINATION.** Crown 8vo. Price 3s. 6d. net.

**MEDICAL CONDUCT AND PRACTICE.** A Guide to the Ethics of Medicine. Crown 8vo. Price 6s. net.

**A MANUAL OF MEDICAL JURISPRUDENCE AND TOXICOLOGY.** Fourth Edition, Crown 8vo., with 38 Illustrations. Price 12s. 6d. net.

**A MANUAL OF PUBLIC HEALTH.** Fourth Edition, containing 25 Illustrations in the text. Crown 8vo. Price 8s. 6d. net.

**MEDICAL HISTORY MANUALS.** Each with 8 page Illustrations. Large Crown 8vo. 4 Volumes now ready. List on application. Price 6s. and 7s. 6d. net.

## THE LAWS OF HEALTH FOR SCHOOLS

By **A. M. MALCOLMSON**, M.D. Containing 35 Illustrations. Crown 8vo. Price 2s. 6d.

## RADIOGRAPHY AND RADIOTHERAPEUTICS

By **ROBERT KNOX**, M.D. Edin., C.M., Edin., M.R.C.S. Eng., L.R.C.P. Lond. Third Edition, Revised and Enlarged. 90 full-page Plates and over 400 Illustrations in the text

Vol. I.—Radiography 35/- net  
" II.—Radiotherapeutics 18s. net

## A TEXT-BOOK OF MIDWIFERY

By **R. W. JOHNSTONE**, M.A., M.D., F.R.C.S.E., M.R.C.P.E. Third Edition, containing 262 Illustrations (3 in colour). Crown 8vo. Price 15s. net.

## GENERAL PRACTICE AND X-RAYS

By **ALICE VANCE KNOX**, M.B., B.Ch. With Chapters on Instrumentation by **ROBERT KNOX**, M.D., C.M., M.R.C.S., L.R.C.P. Crown 8vo., many Illustrations. 10s. 6d. net.

## TUBERCULOSIS OF BONES AND JOINTS IN CHILDREN

By **JOHN FRASER**, M.D., F.R.C.S.E., Ch.M. Containing 51 full-page Plates (2 of them in colour) and 164 Illustrations in the text. Super Royal 8vo. Price 20s. net.

## THE STRUCTURE OF THE FOWL

By **O. CHARNOCK BRADLEY**, M.D., D.Sc., M.R.C.V.S. Containing 73 Illustrations. Crown 8vo. Price 6s. net.

By **JAMES BURNET**, M.A., M.D., M.R.C.P.E. Foolscap 16mo (4½ by 2½). **A DICTIONARY OF MEDICAL TREATMENT AND DIET.** New Edition, Revised. Crown 8vo. Price 5s. net.

**THE POCKET PRESCRIBER.** Fifth Edition. Price 2s. net.

**THE POCKET CLINICAL GUIDE.** Third Edition. Price 2s. 6d. net.

**BLACK'S MEDICAL DICTIONARY.** Edited by **JOHN D. COMRIE**, M.A., B.Sc., M.D., F.R.C.P.E. Seventh Edition, completing 50,000. Revised and much enlarged. Containing 431 Illustrations, 2 being full-page in colour. Demy 8vo. Price 15s. net.

**A. & C. BLACK, LTD., 4, 5, & 6 Soho Square, London, W.1**

**TESTOGAN**

For Men

**THELYGAN**

For Women

*Formula of Dr. Iwan Bloch.**After eight years' clinical experience these products stand as proven specifics.***Indicated in Sexual Impotence and Insufficiency of the Sexual Hormones.**They contain **SEXUAL HORMONES**, i.e., the hormones of the reproductive glands and of the glands of internal secretion.*Special Indications for Testogan:—Sexual infantilism and eunuchoidism in the male. Impotence and sexual weakness.**Climacterium virile. Neurasthenia, hypochondria.**Special Indications for Thelygan:—Infantile sterility. Under-developed mammae, etc. Frigidity. Sexual disturbances in obesity and other metabolic disorders. Climacteric symptoms, amenorrhea, neurasthenia, hypochondria, dysmenorrhea.**Furnished in Tablets for internal use, and in Ampoules, for intragluteal injection.***EXTENSIVE LITERATURE ON REQUEST TO THE  
CAVENDISH CHEMICAL CORPORATION,****Empire House, 175 Piccadilly, London, W.1.***English Distributors:—BUTLER & CRISPE, LONDON, E.C.1.  
MESSRS. MAY, ROBERTS & Co., Ltd., Dublin, Plymouth & London.***For INDIGESTION  
and DYSPEPSIA****Lactopeptine****LACTOPEPTINE IS A NON-SECRET  
REMEDY WITH ITS FORMULA  
ON THE LABEL OF EACH BOTTLE.**

**T**HE digestive ferments (pepsin, pancreatine, diastase) used in the manufacture of Lactopeptine are of the highest grade. Although this preparation has been prescribed by the Medical Profession for over 40 years, its ingredients have been constantly improved and kept up to date by the most advanced pharmaceutical practice.

When prescribing specify in all cases Lactopeptine (Richards.) Dispensed in 1 oz. (4/6) and  $\frac{1}{2}$  oz. (2/9) bottles, in powder or tablet form. To prevent substitution, Medical men are requested to prescribe Lactopeptine in original bottles.

*Samples (powder and tablet form) free  
of charge on receipt of professional card.***JOHN MORGAN RICHARDS & SONS, LTD.,****46 HOLBORN VIADUCT, LONDON, E.C.1.**



# SALT & SON, LTD.,

HEAD OFFICES:—

6 CHERRY ST., BIRMINGHAM.

LONDON CONSULTING ROOMS:

14-18 BLOOMSBURY STREET, W.C.

MUSEUM 3845. Fully qualified Nurse in attendance.


BY APPOINTMENT ONLY.

*Orthopædic Appliance Specialists.*


*Artificial Limb Manufacturers.*

## Salts Gastropnoxis Belt (Regd. No. 682303 P. Pat 11098/21.)

as designed for JOHN F. HALL-EDWARDS. Major (late R.A.M.C.)  
L.R.C.P. Edin., D.M.R. & E. Contab., F.R.S. Edin.



FRONT VIEW.



SIDE VIEW

Write for Particulars and Measurement Forms.

Extract from article "The Diagnosis and Treatment of Gastropnoxis," by Major Hall-Edwards, in B.M.J., May 14, 1921:—

"By far the most important item in the treatment is the special abdominal belt for holding the stomach up."

SEND FOR CATALOGUES OF

BELTS AND CORSETS. ELASTIC HOSIERY.

TRUSSES.

ARTIFICIAL LIMBS.

and DEFORMITY APPARATUS.

Hæmorrhoidal

**ANUSOL**  
Regd

Suppositories

**Anusol Suppositories** give prompt relief from pain, and make possible easy natural evacuations.

**Anusol Suppositories** cleanse, disinfect and heal inflamed membrane.

**Anusol Suppositories** are entirely free from narcotics and caustic ingredients. They dissolve slowly without giving rise to pain, irritation or leakage.

*The formula of Anusol Suppositories appears on each Box.*

CHAS. ZIMMERMANN & CO. (CHEM.), LTD.,  
9-10 St. Mary-at-Hill, London, E.C.3.

# Antikamnia Tablets



## For Influenza and La Grippe.



For the headache pain and general soreness give a five-grain Antikamnia Tablet crushed with a little water; if the pain is very severe, two tablets should be given. Repeat every two or three hours as required. One single ten-grain dose is often followed by complete relief.

## Laryngeal Cough



Frequently remains after an attack of Influenza, and has been found stubborn to yield to treatment. There is an irritation of the larynx, huskiness, and a dry and wheezing cough, usually worse at night. The prolonged and intense paroxysms of coughing are controlled by **ANTIKAMNIA and CODEINE TABLETS**, and with the cessation of the coughing, the laryngeal irritation subsides. Antikamnia Tablets are the least depressing of all the drugs that can exercise so extensive a control of pain, and also least disturbing to the digestive and other organic functions.

**Analgesic. Antipyretic. Anodyne.**

Antikamnia Preparations in 1-oz. packages only.

JOHN MORGAN RICHARDS & SONS, LTD.  
46-47 Holborn Viaduct, LONDON, E.C.1



DR. JUNIOR: "What did you think of Purdy's paper—I suppose you agree with him that Antiphlogistine is the ideal poultice?"

DR. SENIOR: "Antiphlogistine, my young colleague and friend, is vastly more than a mere poultice."

DR. JUNIOR: "You think it replaces the sour-smelling bread-and-milk affair, or the slimy, quickly cooling linseed-meal nuisance——"

DR. SENIOR: "I no longer use those bacteria-breeding cataplasms. To say nothing of the personal appeal of Antiphlogistine, it is physiologically active from the first contact; it is physically clean, convenient and agreeable to the most fastidious patient——"

DR. JUNIOR: "And always available at any drug store."

DR. SENIOR: "Exactly—Antiphlogistine may safely be called our most scientifically satisfactory method of applying moist heat to an inflamed part and maintaining it. Most patients, when it is properly applied, either soon go to sleep, or say how comfortable it is."

DR. JUNIOR: "I suppose I can get a reprint of Dr. Purdy's Paper?"

DR. SENIOR: "The Antiphlogistine company will probably have excerpts of the article."



# Valentine's Meat-Juice

In **Diarrhoea, Dysentery and Cholera Infantum** where it is **Essential to Conserve the Weakened Vital Forces** without **Irritating the Digestive Organs, Valentine's Meat-Juice** demonstrates its **Ease of Assimilation** and **Power to Sustain and Strengthen.**

## Diseases of Children

**W. T. Watt, M. D., Director Imperial Medical College, Tientsin, China:** "In cases of Infantile Diarrhoea, which weakens and debilitates a child rapidly, I have found Valentine's Meat-Juice a great stimulant and quick restorative of vitality. Three years ago, when an epidemic broke out in Tientsin, I ordered my staff to try your Meat-Juice, which justified all expectations, having been satisfactory to patients and physicians alike."

**Henry N. Read, M. D., Prof Diseases of Children, Long Island College Hospital, Brooklyn, N. Y.:** "I have long used Valentine's Meat-Juice, especially in Diseases of Children, and esteem it highly. It has proven most valuable in the Enteric Disorders of Children, both in my practice and Hospital work."



---

For Sale by European and American Chemists and Druggists

---

**VALENTINE'S MEAT-JUICE COMPANY.**

**Richmond, Virginia, U. S. A.**